

# Evidence for Genes on Chromosome 2 Contributing to Alcohol Dependence With Conduct Disorder and Suicide Attempts

Danielle M. Dick,<sup>1\*</sup> Jacquelyn Meyers,<sup>1</sup> Fazil Aliev,<sup>1</sup> John Nurnberger Jr.,<sup>2</sup> John Kramer,<sup>3</sup> Sam Kuperman,<sup>3</sup> Bernice Porjesz,<sup>4</sup> Jay Tischfield,<sup>5</sup> Howard J. Edenberg,<sup>2</sup> Tatiana Foroud,<sup>2</sup> Marc Schuckit,<sup>6</sup> Alison Goate,<sup>7</sup> Victor Hesselbrock,<sup>8</sup> and Laura Bierut<sup>7</sup>

<sup>1</sup>Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, Richmond, Virginia

<sup>2</sup>Indiana University School of Medicine, Indianapolis, Indiana

<sup>3</sup>University of Iowa College of Medicine, Iowa City, Iowa

<sup>4</sup>SUNY Health Science Center at Brooklyn, Brooklyn, New York

<sup>5</sup>Rutgers University, Piscataway, New Jersey

<sup>6</sup>University of California, San Diego VA Medical Center, San Diego, California

<sup>7</sup>Washington University in St. Louis, St. Louis, Missouri

<sup>8</sup>University of Connecticut Health Center, Farmington, Connecticut

Received 27 August 2009; Accepted 22 February 2010

Twin studies provide strong evidence that there is a shared genetic liability that predisposes to a number of different psychiatric outcomes related to behavioral disinhibition. Further, alcohol dependence comorbid with other disinhibitory disorders is particularly heritable. Chromosome 2p14-2q14.3 has been linked to multiple psychiatric conditions related to behavioral undercontrol. In the Collaborative Study on the Genetics of Alcoholism (COGA), we previously reported linkage to this region with alcohol dependence (AD), suicide attempts (SUI), and conduct disorder (CD). In this study, we follow-up on these previous reports of linkage by combining the phenotypes in analyses that jointly consider the presence of multiple conditions. Linkage analyses of the combined phenotype of AD with CD or SUI results in a maximum LOD score of 5.4 in this region. In addition to this primary linkage peak, independent samples have reported linkage to other alcohol-related phenotypes across chromosome 2. Accordingly, we followed-up these linkage signals by testing for association with SNPs across chromosome 2 in a case-control sample, in which a subset of the cases consisted of alcohol-dependent probands from the linkage sample. We find evidence of association with the combined AD with CD or SUI phenotype, with 23 genes surviving permutation testing. The number of associated genes across the chromosome may explain the persistent linkage findings reported on chromosome 2 across a number of independent studies of alcohol and disinhibitory phenotypes. Further, none of the genes were located directly under the primary COGA linkage peak, which has implications for association tests following-up linkage peaks. © 2010 Wiley-Liss, Inc.

## How to Cite this Article:

Dick DM, Meyers J, Aliev F, Nurnberger J Jr, Kramer J, Kuperman S, Porjesz B, Tischfield J, Edenberg HJ, Foroud T, Schuckit M, Goate A, Hesselbrock V, Bierut L. 2010. Evidence for Genes on Chromosome 2 Contributing to Alcohol Dependence With Conduct Disorder and Suicide Attempts.

Am J Med Genet Part B 153B:1179–1188.

**Key words:** alcoholism; genetics; linkage; association; behavioral disinhibition

Additional Supporting Information may be found in the online version of this article.

Grant sponsor: NARSAD; Grant numbers: U10AA008401, N01-HG-65403; Grant sponsor: National Institute on Alcohol Abuse and Alcoholism (NIAAA); Grant sponsor: National Institute on Drug Abuse (NIDA).

\*Correspondence to:

Dr. Danielle M. Dick, Ph.D., Department of Psychiatry, Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, PO Box 980126, Richmond, VA 23298-0126.

E-mail: ddick@vcu.edu

Published online 16 April 2010 in Wiley Online Library (wileyonlinelibrary.com).

DOI 10.1002/ajmg.b.31089

## INTRODUCTION

Twin studies provide strong evidence of shared genetic liability across a number of different psychiatric conditions. For example, in a large study of the genetic architecture of the major common psychiatric and substance use disorders, Kendler et al. [2003], using data from the Virginia Twin Registry, demonstrated that there were two broad genetic factors: one that contributed to externalizing disorders (alcohol dependence, drug abuse/dependence, childhood conduct disorder, and adult antisocial behavior) and a second that contributed to internalizing disorders (major depression, generalized anxiety disorder and phobia). These findings have implications for gene identification efforts, as they suggest that some genes may not be specific to any one disorder, but rather, may predispose to a variety of psychiatric outcomes. Furthermore, individuals meeting a psychiatric diagnosis are often a heterogeneous group clinically. For example, in the case of alcohol dependence, affected individuals often vary on a number of important dimensions, including age of onset, course of illness, and the presence of comorbid conditions [Cloninger, 1987; Babor et al., 1992; Hesselbrock and Hesselbrock, 1994; Finn et al., 1997]. Evidence from twin and family studies suggests that alcohol dependence with comorbid disinhibitory disorders may represent a more heritable form of the disorder [Pickens et al., 1991, 1995; Johnson et al., 1996; Ohannessian et al., 2004] suggesting that comorbid, or combined, phenotypes may be particularly relevant for gene identification efforts.

Data from the Collaborative Study on the Genetics of Alcoholism (COGA) suggest that chromosome 2p14-2q14.3 may contain a gene (or genes) with pleiotropic effects on alcohol dependence and related psychiatric conditions. Initially, linkage was detected near the marker D2S379 (LOD = 3.0) with an alcohol dependence (AD) phenotype, defined as meeting diagnostic criteria according to the DSM-III-R and Feighner classification systems [Foroud et al., 2000]. Subsequently, linkages to the same region were identified with the phenotypes of suicide attempts (SUI) [Hesselbrock et al., 2004] and conduct disorder (CD) [Dick et al., 2003]. These findings are robust, with replication reported in multiple independent samples: Suicide attempts were linked to this same region of chromosome 2 in pedigrees affected with early-onset major depression [Zubenko et al., 2004] and with bipolar disorder [Willour et al., 2007]. Linkage of conduct disorder to this region was replicated in the Irish Affected Sib Pair Study for Alcohol Dependence [Kendler et al., 2006a]. Further analysis of a subset of the COGA pedigrees on whom genome-wide SNP linkage data were produced for the Genetic Analysis Workshop 14 (GAW14) [Edenberg et al., 2005] suggested that the chromosome 2 linkage finding for alcohol dependence was one of the most robust linkage signals in the sample [Doan et al., 2005; Wang et al., 2005; Wiener et al., 2005]. Extension of the linkage markers available at the ends of the chromosomes in the GAW14 SNP set also suggested evidence for linkage peaks with alcohol dependence at the p and q ends of the chromosome, in addition to the primary centromeric linkage peak discussed here [Wang et al., 2005; Wiener et al., 2005; Agrawal et al., 2008]. Linkages to alcohol dependence [Wilhelmsen et al., 2005] and the related traits of alcohol withdrawal [Kuo et al., 2006] and

smoking [Straub et al., 1999; Goode et al., 2003] have also been reported on chromosome 2 in independent samples. Although linkage to a comorbid habitual smoking and alcohol dependence phenotype has been reported to chromosome 2p in COGA [Bierut et al., 2004], the finding is largely due to the alcohol dependence phenotype.

The phenotypes of alcohol dependence, conduct disorder, and suicide attempts, which show linkage on chromosome 2, are all characterized by elements of impulsivity and behavioral undercontrol. Conduct disorder is a robust predictor of both concurrent and future alcohol problems [Crowley et al., 1998; Moss and Lynch, 2001; White et al., 2001]. Furthermore, numerous twin studies indicate that the overlap between childhood conduct disorder and adult alcohol dependence is largely due to shared genetic factors [Slutske et al., 1998; Krueger, 1999; Young et al., 2000; Kendler et al., 2003]. This common genetic liability is thought to be a predisposition toward behavioral undercontrol/disinhibition, which can manifest as conduct disorder in childhood and alcohol dependence later in life [Slutske et al., 2002]. It has been demonstrated that *GABRA2*, a gene associated with alcohol dependence in adults [Edenberg et al., 2004; Lappalainen et al., 2005; Enoch, 2008; Soyka et al., 2008] is associated with CD symptoms and externalizing behavior in adolescents [Dick et al., 2006, 2009], providing evidence that variations in one gene can manifest as different conditions at different stages of the life cycle. Electrophysiological endophenotypes, which are thought to index genetic vulnerability to psychiatric phenotypes, are also shared across substance use disorders and conduct disorder [Iacono et al., 1999; Porjesz et al., 2005]. For example, a reduced P3 event-related potential amplitude has been found among adolescents with both substance use disorders and externalizing disorders [Iacono et al., 2002]. Suicide attempts are also considerably elevated in individuals with alcohol dependence and conduct disorder [Kessler et al., 1999]. A recent study by Conner et al. [2009], found that proactive aggression (unemotional aggression executed for reward) is associated with both suicide attempts and suicidal ideation among inpatient substance abuse patients. Furthermore, in the COGA adult sample, unplanned suicide attempts among persons with alcohol dependence are associated with externalizing behaviors, including antisocial behavior and alcohol-related aggression [Conner et al., 2007]. Suicidal behavior is influenced by genetic factors [Bondy et al., 2006], and it is thought that the predisposition to suicidal behavior reflects a heritable liability to personality traits such as impulsivity and aggression [Baud, 2005]. To the extent that these characteristics also predispose to substance use and externalizing problems, it is reasonable to hypothesize that suicidal behavior may represent another manifestation of an underlying predisposition toward behavioral disinhibition. Longitudinal studies have found that behavioral disinhibition measured in childhood/adolescence predicts both the development of substance use disorders and a propensity toward suicidal behavior in young adulthood [Tarter et al., 2004].

Additional follow-up of the linkage signals observed across the three phenotypes in COGA suggested that it was not simply the same individuals contributing to the results across the phenotypes; for example, only half of the individuals who had attempted suicide also had alcohol dependence, and only 25% of the individuals with

alcohol dependence have childhood conduct disorder. To the extent that these phenotypes represent an underlying liability to behavioral disinhibition, considering these phenotypes jointly in linkage analyses, rather than analyzing each individually, should enhance the power to detect linkage if there is a gene (or genes) in the region that is predisposing to this broad constellation of psychiatric outcomes. Further, to the extent that AD characterized by suicide attempts and conduct disorder may represent a more homogeneous and heritable form of the disorder, considering a combined phenotype would also increase our power to detect linkage.

Here we report a series of linkage and association analyses in the COGA sample to follow up the individual reports of linkage to alcohol dependence, conduct disorder, and suicide attempts previously reported on chromosome 2. First, we conducted linkage analyses that jointly consider the previously linked phenotypes in the COGA sample. We conducted these analyses using the high risk COGA family-based sample, in which both microsatellite [Reich, 1996] and SNP [Edenberg et al., 2005] linkage panels are available. Secondly, we tested for association with SNPs across chromosome 2 in a case-control sample, which included a subset of the alcohol-dependent probands from the linkage sample.

## METHODS

### Sample

COGA is a multi-site project, with the goal of identifying genes contributing to alcoholism and related phenotypes. Proband were identified through inpatient or outpatient alcohol treatment programs at six sites around the United States and were invited to participate if they had a sufficiently large family (usually sibships >3 with parents available) with two or more members in a COGA catchment area [Begleiter et al., 1995]. The institutional review boards of all participating centers approved the study. Written consent was obtained from all study participants. Additional details about the study have been published previously [Reich et al., 1998; Foroud et al., 2000; Edenberg et al., 2004]. All individuals aged 18 or older were interviewed using the Semi-Structured Assessment for the Genetics of Alcoholism (SSAGA) [Bucholz et al., 1994; Hesselbrock et al., 1999]. Alcoholism was defined by the presence of a DSM-III-R alcohol dependence diagnosis [Diagnostic and Statistical Manual of Mental Disorders: III-R, 1987], plus definite alcoholism according to Feighner Criteria [1972]. The SSAGA makes a diagnosis of childhood conduct disorder according to DSM-III-R through retrospective report of behavioral problems evidenced before the age of 15; diagnoses required the presence of three or more symptoms. Suicide attempts were assessed using an item in the SSAGA that asks individuals if they have ever tried to kill themselves.

### Molecular Methods and Analysis

**Linkage samples and analyses.** A microsatellite linkage scan was conducted on a sample of 2,282 individuals from 262 families densely affected with alcohol dependence [Reich et al., 1998; Foroud et al., 2000]. Genotyping for the microsatellite linkage scan was carried out in laboratories at Indiana University and Wash-

ington University in St. Louis using radioactive and fluorescence-based detection systems, as described previously [Reich et al., 1998; Foroud et al., 2000]. The current analyses are based on a map of 315 autosomal microsatellite markers with an average intermarker distance of 11.5 cM. Pedigrees were checked for non-Mendelian inheritance using the GeneMaster database and the programs CRIMAP [Green, 1990] and USERM13 [Boehnke, 1991]. Recombination-based marker maps were generated from the sample using CRIMAP. Maximum likelihood estimates of marker allele frequencies were computed from the data using USERM13. The majority of the sample (84%) was Caucasian; 13% was African American, and <3% reported mixed or other ethnicities. Alleles were coded separately for Caucasians and African Americans/others to take into account allele frequency differences between the populations.

In addition to the primary COGA linkage sample, a subset of the COGA linkage families were selected for additional genotyping as part of GAW14 [Edenberg et al., 2005]. A subset of 1,364 individuals from 143 families was genotyped on an Illumina panel of 4,596 SNPs intended as a linkage SNP panel. Parallel to the full COGA sample, the majority of the sample was Caucasian (83%), with 13% African American, and 4% other. Because linkage disequilibrium can produce spurious inflations in identity-by-descent (IBD) estimates and inflate information content as a result [Huang et al., 2004], we conducted linkage on a thinned panel of 1,717 SNPs, in which all SNPs with  $r^2 \geq 0.1$  with any other SNP within 1 Mb were deleted [Agrawal et al., 2008]. The thinned map provided similar information content across the genome when compared to the full panel of SNPs [Hinrichs et al., 2005].

Non-parametric, multipoint methods of linkage analysis for affected sibling pairs were employed, first using the microsatellite linkage panel, and subsequently with the SNP linkage panel, using the program ASPEX (Hinds & Risch, 1999). The linkage analyses were performed using only those affected siblings with both parents genotyped (sib\_ibd), which allows unambiguous estimation of IBD. This type of analysis results in greater accuracy in the estimate of marker allele sharing among affected siblings. Analyses were performed using all possible pairs of affected siblings [ $n(n-1)/2$ ], where  $n$  = number of affected siblings in a nuclear family]. First, each of the phenotypes (alcohol dependence, conduct disorder, and suicide attempts) was analyzed individually, parallel to the previous reports (however, we note that the results are slightly different than the original publications, reflecting an updated genetic map). Subsequently, all three phenotypes were combined, such that affected status was defined by the presence of alcohol dependence (AD) or conduct disorder (CD) or suicide attempts (SUI). Finally, an analysis focused on only the subset of alcohol-dependent individuals who also have either conduct disorder or a suicide attempt (AD with CD or SUI) was conducted.

**Association analyses.** SNP data were available across chromosome 2, generated as part of the genome-wide association study (GWAS) of the COGA sample by the Center for Inherited Disease Research, using the Illumina HumanHap1M BeadChip platform, and are available through dbGaP. After all data cleaning, genome wide data were available for 1905 individuals [Edenberg et al., 2010]. All 1,205 cases met criteria for DSM-IV Alcohol

Dependence, as assessed by the SSAGA, and all 700 controls were screened against Alcohol Dependence and related substance use disorders. Three hundred twenty-four of the cases were drawn from families included in the primary COGA linkage sample. Males were overrepresented in the cases. Additional details about the COGA GWAS sample are available in Edenberg et al. [2010].

Because all cases genotyped in the GWAS sample were affected with AD, the association analyses focused on the combined AD with CD or SUI phenotype. Individuals were considered affected if they met criteria for AD and either CD or SUI, in addition to the primary AD case phenotype: 511 individuals (of the original 1,205 cases) met criteria for the AD with CD or SUI phenotype. A subset of individuals in the COGA GWAS sample ( $N = 321$ ) were not assessed for suicide attempts; accordingly, when unaffected by conduct disorder, their combined "case" status was unknown, so 231 individuals were excluded from analyses for this reason. Additionally, 40 controls were excluded from analyses because of previous suicide attempts ( $N = 18$ ) or the presence of a conduct disorder diagnosis ( $N = 22$ ). Of the 511 AD cases who met criteria for the AD with either CD or SUI phenotype, 218 (42.7%) had a reported suicide attempt and 367 (71.8%) met criteria for conduct disorder; 74 individuals (14.5%) met criteria for both. 70.8% were male and 29.2% female. The age range was 18–74 (mean = 39.41). Of this group, 74.0% were European American ( $n = 378$ ), 24.9% were African American ( $n = 127$ ), and 1.2% were of another ethnicity ( $n = 6$ ).

The program Plink [Purcell et al., 2007] (<http://pngu.mgh.harvard.edu/purcell/plink/>) was used to conduct all association tests. Logistic regression association analyses using an additive genetic model were conducted on the combined phenotype defined by the presence of AD *with* CD or SUI. The association analyses were run on each of the 82,562 SNPs genotyped across chromosome 2, using as covariates sex and ethnicity, as defined based on principal component-based analysis performed in PLINK to cluster the samples along with HapMap reference samples. Additional details are available in [Edenberg et al., 2010]. There were 1,134 genes across the chromosome. We used a gene-based strategy for association testing, an approach which can offer the advantage of identifying association to genes where multiple common variants may exist, rather than focusing on a single SNP [Moskvina et al., 2009]. In previous studies of AD in the COGA sample, we have frequently observed multiple signals across the gene [e.g., Wang et al., 2004;

Edenberg et al., 2007; Dick et al., 2008; Wetherill et al., 2008]. The observation of multiple (not entirely correlated) signals across a gene enhances our confidence that an observed association is real.

Permutation tests were conducted for all genes yielding at least 1  $P$ -value  $< 0.001$ , as well as for genes which yielded at least two SNPs with  $P$ -values  $< 0.01$ , with an  $r^2 < 0.8$  between those SNPs. This approach captured genes with single associated SNPs or genes with converging evidence for association from multiple SNPs. To take into account the LD structure across SNPs in a gene, Plink begins with the most significant SNP in the gene, removes SNPs with  $r^2 > 0.80$ , and continues this process for all SNPs with  $P < 0.05$  to obtain an independent group of associated SNPs. The  $P$ -value associated with the average test statistic across the remaining SNPs is compared across 10,000 permutation runs, yielding an empirical  $P$ -value for the gene based on the independent, significant SNPs observed in the gene.

## RESULTS

### Linkage Analyses

Table I shows the number of affected sibling pairs, LOD scores, and allele sharing for each of the phenotypes using the primary COGA microsatellite linkage panel and the SNP linkage panel available on a subset of the full COGA sample. Since LOD scores are influenced by sample size, and different numbers of sibling pairs were available for each of the phenotypes (and between the microsatellite and SNP linkage panels), allele sharing is provided in Table I to allow comparisons across the phenotypes and across the two sets of linkage analyses. Although the LOD scores differ across the microsatellite and SNP linkage panels (not surprising due to differences in sample sizes and methodology), the pattern of allele sharing observed using the microsatellite and SNP linkage panels is largely consistent. Of the individual phenotypes, CD and SUI yield higher rates of allele sharing than AD. The allele sharing for AD *with* CD or SUI was considerably elevated compared to the full AD analysis, and the AD *or* CD or SUI showed only a very small elevation in allele sharing. Further, the combined phenotype (AD *with* CD or SUI) showed elevated allele sharing compared to the broader AD *or* CD or SUI phenotype. This suggests, across both sets of analyses, that the phenotype of AD *with* CD or SUI represents a more genetically homogeneous subgroup influenced by gene(s) in the region. This was the phenotype used in subsequent association testing.

**TABLE I. Results From Affected Sibling Pair Linkage Analyses Using Microsatellite Markers and SNP Linkage Panel**

Phenotype	Microsatellite markers				SNPs			
	# Sibling pairs	Maximum LOD Score	Position [cM]	Allele-sharing (%)	# Sibling pairs	Maximum LOD Score	Position [cM]	Allele-sharing (%)
Alcohol dependence [AD]	797	2.9	114	55	384	1.03	117	57
Conduct disorder [CD]	113	2.4	117	64	76	3.7	102	86
Suicide attempts [SUI]	58	2.7	117	68	40	2.1	99	84
AD or CD or SUI	988	3.7	114	56	531	1.5	132	57
AD with CD or SUI	128	3.6	113	66	90	5.4	109	77

TABLE II. Genes on Chromosome 2 Yielding Evidence for Association at  $P < 0.05$  After Permutation Testing

GENE symbol	Gene name	BP position	NSNP	NSIG	ISIG	EMP	Significant SNPs
NTSR2	Neurotensin receptor 2	11724667	3	2	2	0.0055	rs12612207 rs4669765
TRIB2	Tribbles homolog 2	12784955	12	3	3	0.0221	rs890069 rs10189072 rs17465002
PPM1G	Protein phosphatase 1G, magnesium-dependent, gamma isoform	27459602	18	1	1	0.0003	rs2384629
MEMO1	Mediator of cell motility 1	31964229	65	3	3	0.0216	s17011668 rs17011667 rs3769609
HAAO	3-Hydroxyanthranilate 3,4-dioxygenase	42849352	13	7	6	0.0327	rs3755541 rs13027051 rs2374442 rs3816184 rs3816182 rs737148
MTIF2	Mitochondrial translational initiation factor 2	55345558	17	2	2	0.0171	rs6721728 rs6707902
CCDC139	Pseudouridylylate synthase 10	61068848	33	5	5	0.0298	rs7564317 rs6708713 rs6715485 rs11691111 rs9309333
EHPB1	EH domain binding protein 1	62977118	146	22	10	0.0246	rs2710638 rs13006926 rs4671453 rs1123508 rs2018650 rs13027462 rs1468748 rs4671052 rs7557501 rs17432497
BUB1	Budding uninhibited by benzimidazoles 1 homolog	111135899	16	5	3	0.0240	rs7609252 rs13398617 rs12053209
TTL	Tubulin tyrosine ligase	112962482	25	7	7	0.0302	rs6726169 rs6718489 rs1561266 rs7570679 rs7578685 rs34179430 rs7559710
CKAP2L	Cytoskeleton associated protein 2-like	113230296	12	3	3	0.0244	rs6731822 rs10209160 rs7577241
MGAT5	Mannosyl (alpha-1,6-)-glycoprotein beta-1,6-N-acetylglucosaminyltransferase	134754387	96	4	4	0.0318	rs16830319 rs7564658 rs11887041 rs3791269
ARHGAP15	Rho GTPase activating protein 15	143831058	212	13	13	0.0344	rs6430025 rs10205708 rs6714864 rs11890035 rs2890714 rs10172663 rs13406291 rs17814868 rs17229439 rs16858636 rs11681284 rs10195682 rs6704667
KIAA1189	Ermin, ERM-like protein	157883898	7	1	1	0.0006	rs17282140
COBLL1	COBL-like 1	165344115	63	13	13	0.0106	rs355895 rs355907 rs355865 rs355846 rs355868 rs355900 rs355844 rs6414069 rs355825 rs355810 rs6748091 rs355911 rs355849
FAM130A2	Cysteine-serine-rich nuclear protein 3	166211286	39	2	2	0.0290	rs6720974 rs17251144
LPR2	LiPocalin-related protein 2	169733586	118	10	9	0.0416	rs2239602 rs2302694 rs4140872 rs13417486 rs2075252 rs990626 rs2239591 rs2239596 rs1548936
CHN1	Chimerin 1	175484629	84	4	4	0.0438	rs16862927 rs1193623 rs1193630 rs2605285
PRKRA	Protein kinase, interferon-inducible double stranded RNA-dependent activator	179015447	8	2	2	0.0112	rs13427914 rs13392094

(Continued)

TABLE II. (Continued)

GENE symbol	Gene name	BP position	NSNP	NSIG	ISIG	EMP	Significant SNPs
PPP1R1C	Protein phosphatase 1, regulatory (inhibitor) subunit 1C	182688766	59	6	5	0.0210	rs10451546 rs16822592 rs1882212 rs16822590 rs16867518
LOC402117	Von Willebrand factor C domain-containing protein 2-like	215092215	52	18	15	0.0204	rs10932539 rs9288497 rs10932540 rs7558283 rs6719667 rs13425618 rs4629153 rs12612233 rs4321367 rs13417343 rs4673830 rs6760280 rs11904475 rs6729204 rs6740246 rs16858808 rs16858816 rs16858811
IL8RA	Interleukin 8 receptor, alpha	218737177	11	3	3	0.0084	
FARP2	FERM, RhoGEF and pleckstrin domain protein 2	241951484	43	8	6	0.0458	rs3771550 rs3771561 rs2240482 rs6723363 rs1476698 rs764081


The base pair position corresponds to the most significant SNP in each gene; NSNP, the total number of SNPs that were tested across the gene; NSIG, the total number of SNPs that yielded  $P$ -values  $< 0.05$  across the gene; ISIG, the number of independent SNPs yielding  $P$ -values  $< 0.05$  (as determined by Plink); EMP, the empirical  $P$ -value for the gene based on permutation tests.

## Association Analyses of the Phenotype AD With CD or SUI

Sixty-one genes across chromosome 2 yielded at least one SNP with  $P$ -value  $< 0.001$ , or at least two SNPs (with an  $r^2 < 0.8$ ) with  $P$ -values  $< 0.01$ , and were subjected to permutation testing. Twenty-three genes had an empirical  $P < 0.05$ . Table II shows the genes that yielded empirical  $P$ -values  $< 0.05$  in permutation tests. Table II lists the number of SNPs that were tested across each gene, the total number of SNPs that yielded  $P$ -values  $< 0.05$  across the gene, the number of independent SNPs yielding empirical  $P$ -values  $< 0.05$ , the empirical  $P$ -value for the gene based on permutation tests, and the significant SNPs. The original  $P$ -values for those SNPs are presented in Supplemental Table I. In Figure 1, the approximate location of these genes is listed along the top of the LOD score graph (based on the microsatellite linkage analyses from the full COGA family sample) to illustrate roughly their chromosomal positions with respect to the linkage peak. Interestingly, none of these genes was located within a 1 LOD drop of the primary, centromeric linkage peak. Rather, those genes in closest proximity to the linkage peak were clustered on either side of the peak. In addition, consistent with previous reports of linkage across the chromosome, additional genes surviving permutation testing were located across chromosome 2.

## DISCUSSION

In this article we follow-up on evidence for linkage on chromosome 2 with alcohol dependence [Foroud et al., 2000; Wilhelmsen et al., 2005], conduct disorder [Dick et al., 2003; Kendler et al., 2006a], and suicide attempts [Hesselbrock et al., 2004; Zubenko et al., 2004; Willour et al., 2007] in the COGA sample, and independent samples [Zubenko et al., 2004; Kendler et al., 2006b; Willour et al., 2007]. These phenotypes have shown consistent linkage to the centromeric region of chromosome 2 across multiple samples. In addition, there have been linkages reported to alcohol and related phenotypes on the p and q arms of chromosome 2 [Straub et al., 1999; Goode et al., 2003; Wang et al., 2005; Wiener et al., 2005; Wilhelmsen et al., 2005; Kuo et al., 2006; Agrawal et al., 2008]. We find that the phenotype of alcohol dependence with either conduct disorder or suicide attempts results in a maximum LOD score of 5.4 in the centromeric region of chromosome 2. Although the joint phenotype of AD or CD or SUI produced a similar LOD score in the microsatellite linkage analyses, the allele sharing was considerably lower, and in the SNP linkage scan, the LOD score was considerably lower. One hypothesis for this pattern of results is that limiting analyses to the AD individuals *with* CD or SUI yielded a more homogeneous group with a more heritable, behaviorally disinhibited phenotype. Not only is AD more narrowly defined by this joint phenotype, so might be SUI. By requiring at least some of the suicide attempts to be comorbid with alcohol problems, we may have largely retained "externalizing" attempts and eliminated attempts that reflected planned, internalizing-based behavior. The subset of AD individuals in the association sample who also met criteria for conduct disorder or a suicide attempt were a more severe subset of the cases by a number of criteria: they were more likely to meet criteria for an illicit drug dependence diagnosis (77.1% vs. 51.5%,  $P < 0.001$ ), had



**FIG. 1.** Lod score graphs for each of the phenotypes using the microsatellite linkage panel across chromosome 2. Genes with empirical  $P < 0.05$  are shown according to their approximate position along the top of the graph. [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]

a higher mean number of illicit drug dependence symptoms (12.10,  $SD = 10.10$  in AD cases with CD/SUI; 6.53,  $SD = 7.74$  in other AD cases;  $P < 0.001$ ), had a younger age of onset of AD (23.86,  $SD = 7.37$  in AD cases with CD/SUI; 28.12,  $SD = 9.08$  in other AD cases;  $P < 0.001$ ), were more likely to meet criteria for alcohol withdrawal (59.3% vs. 51.3%,  $P = 0.01$ ), and had a slightly higher mean symptom count for alcohol dependence symptoms (5.72 vs. 5.27;  $P < 0.001$ ). Several of the previous reports of associated genes in the COGA sample have been driven by the more severe, comorbid AD individuals in the sample [Edenberg et al., 2006; Dick et al., 2007; Edenberg et al., 2007; Wetherill et al., 2008].

The availability of SNP data across chromosome 2 as part of the GWAS panel allowed us the opportunity to follow-up the chromosome 2 findings without having to a priori define a targeted region. We previously conducted a systematic SNP screen of LD-tagging SNPs across a 2 LOD support interval bracketing a linkage peak on chromosome 7 in the COGA sample [Dick et al., 2008]. In that systematic screen, strong evidence for association was detected with only a single gene (*ACN9*), a surprising result since we know that linkage is not a powerful technique for detecting genes of small effect, and we hypothesize that linkage peaks that are detected for complex traits represent the involvement of multiple genes in the region, an idea that has been supported in following up other linkage peaks in the COGA sample, for example on chromosome 4 [Edenberg et al., 2004, 2006, 2007]. In fact, several other genes have been detected in the vicinity of the chromosome 7 peak in COGA [Hinrichs et al., 2006; Wang et al., 2004, 2007], though these were located outside the 2 LOD support interval. This pattern of results may also reflect the fact that the localization of linkage peaks is known to be imprecise [Roberts et al., 1999]. Accordingly, we made the decision here to take advantage of the existent data across chromosome 2 to test for association. This allowed us to evaluate the location(s) of associated genes with respect to the primary linkage peak. It also allowed us to address the imprecise nature of linkage peaks, the hypothesis of the involvement of multiple genes

in the region (some of which may be missed by narrowly targeting the region just around the peak), and the evidence for linkage to alcohol dependence-related phenotypes across chromosome 2, suggesting there may be relevant susceptibility genes at multiple locations across the chromosome.

The results did prove to be instructive about the localization of linkage peaks with respect to associated genes. Despite the presence of a strong, narrow linkage peak in our sample, none of the genes that passed permutation testing for significance were located with a 1 LOD—or 2 LOD—support interval for the primary, centromeric linkage peak with the AD with CD or SUI phenotype. Although simulations have previously demonstrated that the location of linkage peaks can vary substantially from the position of the underlying variant(s) [Roberts et al., 1999], the empirical results from this study clearly underscore the danger of focusing narrowly on follow-up of linkage regions as defined by the location of the peak. These findings appear to illustrate the hypothetical “worst case scenario” whereby clusters of associated genes are located on either side of the linkage peak, contributing to a peak location in the middle. Because linkage is a within-family test, whereas association is a between-family test, it is also possible that genetic heterogeneity between families could contribute to a detectable linkage signal for which association tests would not be able to detect the underlying genetic variants. This is another possible explanation for the failure to identify any genes directly under the linkage peak. Systematic follow-up across the entirety of chromosome 2 also yields significant evidence of association with multiple other genes that appear to map loosely to the more distal linkage peaks that have been reported on the p and q arms of the chromosome. A preponderance of genes involved in alcohol dependence and related traits across the chromosome likely contributes to the consistent implication of this chromosome in linkage scans.

The original  $P$ -values were more significant than the empirical  $P$ -values (compare Table II and Supplemental Table I) because the permutation tests took into account the multiple SNPs tested across

the genes. Most of the genes that yielded empirical  $P < 0.05$  had multiple independent significant SNPs in the gene, a criteria which COGA has routinely used to bolster confidence in significantly associated genes. However, we note that none of the SNPs would have passed a stringent Bonferroni correction for all SNPs tested across the chromosome, which would have required a  $P$ -value of  $6 \times 10^{-7}$ . Because our study was not completely atheoretical, but rather, we targeted this region of the genome and this particular phenotype based on evidence of linkage in the region, we believe that a gene-based permutation strategy represents a more appropriate means of evaluating significance than the overly conservative Bonferroni correction. However, follow-up in independent samples will ultimately be necessary to evaluate the role of the genes implicated in this study.

None of the genes identified here is currently associated with a large literature involving substance dependence or disinhibitory behavior. One exception may be the gene *NTSR2* that codes for neurotensin receptor 2, a protein that belongs to the G protein-coupled receptor family that activates a phosphatidylinositol-calcium second messenger system. There are previous reports that neurotensin exerts complex effects on the mesolimbic dopamine system that alter motivation and contribute to neuroadaptations associated with psychostimulant drug administration [Garlow et al., 2006; Reynolds et al., 2006]. However, before formulating hypotheses about the potential biological effects of the associated genes, replication in independent samples will be critical.

In conclusion, follow-up of previous independent reports of linkage on chromosome 2 in the COGA sample with alcohol dependence, conduct disorder, and suicide attempts results in a maximal LOD score of 5.4 in the sample with the phenotype of AD with CD or SUI. A systematic screen of SNPs across chromosome 2 with the comorbid AD with CD or SUI phenotype yields evidence of association with 23 genes located across chromosome 2, likely contributing to the preponderance of reported linkages with alcohol dependence and related phenotypes across chromosome 2. These genes may represent novel genes associated with phenotypes resulting from behavioral undercontrol. Confirmation in independent samples will be the next step.

## ACKNOWLEDGMENTS

The Collaborative Study on the Genetics of Alcoholism (COGA), Principal Investigators B. Porjesz, V. Hesselbrock, H. Edenberg, L. Bierut, includes 10 different centers: University of Connecticut (V. Hesselbrock); Indiana University (H.J. Edenberg, J. Nurnberger Jr., T. Foroud); University of Iowa (S. Kuperman, J. Kramer); SUNY Downstate (B. Porjesz); Washington University in St. Louis (L. Bierut, A. Goate, J. Rice, K. Bucholz); University of California at San Diego (M. Schuckit); Howard University (R. Taylor); Rutgers University (J. Tischfield); Southwest Foundation (L. Almasy); and Virginia Commonwealth University (D. Dick). Q. Max Guo is the NIAAA Staff Collaborator. We continue to be inspired by our memories of Henri Begleiter and Theodore Reich, founding PI and Co-PI of COGA, and also owe a debt of gratitude to other past organizers of COGA, including Ting-Kai Li, P. Michael Conneally, and Raymond Crowe, for their critical contributions. This national

collaborative study is supported by the NIH Grant U10AA008401 from the National Institute on Alcohol Abuse and Alcoholism (NIAAA) and the National Institute on Drug Abuse (NIDA). Genotyping services were provided by the Center for Inherited Disease Research (CIDR). CIDR is fully funded through a federal contract from the National Institutes of Health to The Johns Hopkins University, Contract Number N01-HG-65403. This project was supported by a 2007 NARSAD Young Investigator Award to Danielle M. Dick.

## REFERENCES

- Agrawal A, et al. 2008. Linkage scan for quantitative traits identifies new regions of interest for substance dependence in the Collaborative Study on the Genetics of Alcoholism (COGA) sample. *Drug Alcohol Depend* 93(1–2):12–20.
- Babor TF, et al. 1992. Types of alcoholics: Evidence for an empirically derived typology based on indicators of vulnerability and severity. *Arch Gen Psychiatry* 49:599–608.
- Baud P. 2005. Personality traits as intermediary phenotypes in suicidal behavior: Genetic issues. *Am J Med Genet Part C* 133C:34–42.
- Begleiter H, et al. 1995. The Collaborative Study on the genetics of alcoholism. *Alcohol Health Res World* 19:228–236.
- Bierut L, et al. 2004. A genomic scan for habitual smoking in families of alcoholics: Common and specific genetic factors in substance dependence. *Am J Med Genet Part B* 124B:19–27.
- Boehnke M. 1991. Allele frequency estimation from pedigree data. *Am J Hum Genet* 48:22–25.
- Bondy B, Buettner A, Zill P. 2006. Genetics of suicide. *Mol Psychiatry* 11:336–351.
- Bucholz KK, et al. 1994. A new, semi-structured psychiatric interview for use in genetic linkage studies: A report on the reliability of the SSAGA. *J Stud Alcohol* 55(2):149–158.
- Cloninger CR. 1987. Neurogenetic adaptive mechanisms in alcoholism. *Science* 236:410–416.
- Conner KR, et al. 2007. Transitions to, and correlates of, suicidal ideation, plans, and unplanned and planned suicide attempts among 3729 men and women with alcohol dependence. *J Stud Alcohol Drugs* 68(5):654–662.
- Conner KR, Swogger MT, Houston RJ. 2009. A test of the reactive aggression—Suicidal behavior hypothesis: Is there a case for proactive aggression. *J Abnorm Psychol* 118(1):235–240.
- Crowley TJ, et al. 1998. Substance-dependent, conduct-disordered adolescent males: Severity of diagnosis predicts 2-year outcome. *Drug Alcohol Depend* 49:225–237.
- Dick DM, et al. 2003. A genome-wide screen for genes influencing conduct disorder. *Mol Psychiatry* 9:81–86.
- Dick DM, et al. 2006. The role of GABRA2 in risk for conduct disorder and alcohol and drug dependence across developmental stages. *Behav Genet* 36:577–590.
- Dick DM, et al. 2007. Alcohol dependence with comorbid drug dependence: Genetic and phenotypic associations suggest a more severe form of the disorder with stronger genetic contribution to risk. *Addiction* 102(7): 1131–1139.
- Dick DM, et al. 2008. A Systematic single nucleotide polymorphism screen to fine-map alcohol dependence genes on chromosome 7 identifies association with a novel susceptibility gene ACN9. *Biol Psychiatry* 63(11):1047–1053.

- Dick DM, et al. 2009. The role of *GABRA2* in trajectories of externalizing behavior across development and evidence of moderation by parental monitoring. *Arch Gen Psychiatry* 66:649–657.
- Doan BQ, et al. 2005. Application of the propensity score in a covariate-based linkage analysis of the Collaborative Study on the Genetics of Alcoholism. *BMC Genet* 6(Suppl 1): S33.
- Diagnostic and Statistical Manual of Mental Disorders: IIR. 1987. Washington DC: American Psychiatric Association.
- Edenberg HJ, et al. 2004. Variations in *GABRA2*, encoding the  $\alpha 2$  subunit of the GABA-A receptor are associated with alcohol dependence and with brain oscillations. *Am J Hum Genet* 74:705–714.
- Edenberg H, et al. 2005. Description of the data from the Collaborative Study on the Genetics of Alcoholism (COGA) and single-nucleotide polymorphism genotyping for Genetic Analysis Workshop 14. *BMC Genet* 6(Suppl 1): S2.
- Edenberg HJ, et al. 2006. Association of alcohol dehydrogenase genes with alcohol dependence: A comprehensive analysis. *Hum Mol Genet* 15(9):1539–1549.
- Edenberg HJ, et al. 2008. Association of *NFKB1*, which encodes a subunit of the transcription factor NF- $\kappa$ B, with Alcohol Dependence. *Hum Mol Genet* 17(7):963–970.
- Edenberg H, et al. 2010. Genome-wide association study of alcohol dependence implicates a region on chromosome 11. *Alcohol Clin Exp Res* [Epub ahead of print].
- Enoch MA. 2008. The role of GABA(A) receptors in the development of alcoholism. *Pharmacol Biochem Behav* 90(1):95–104.
- Feighner JP, et al. 1972. Diagnostic criteria for use in psychiatric research. *Arch Gen Psychiatry* 26:57–63.
- Finn PR, et al. 1997. Heterogeneity in the families of sons of alcoholics: The impact of familial vulnerability type on offspring characteristics. *J Abnorm Psychol* 106:26–36.
- Foroud T, et al. 2000. Alcoholism susceptibility loci: Confirmation studies in a replicate sample and further mapping. *Alcohol Clin Exp Res* 24: 933–945.
- Garlow SJ, et al. 2006. Genetic analysis of the hypothalamic neurotensin system. *Neuropsychopharmacology* 31(3):535–543.
- Goode EL, et al. 2003. Multiple genome-wide analyses of smoking behavior in the Framingham Heart Study. *BMC Genet* 4(Suppl 1): S102.
- Green PH. 1990. Documentation for CRIMAP, version 2.4.
- Hesselbrock V, Hesselbrock M. 1994. Alcoholism and subtypes of antisocial personality disorder. *Alcohol Alcohol (Suppl 2)*:479–484.
- Hesselbrock M, et al. 1999. A validity study of the SSAGA—A comparison with the SCAN. *Addiction* 94(9):1361–1370.
- Hesselbrock V, et al. 2004. The search for genetic risk factors associated with suicidal behavior. *Alcohol Clin Exp Res* 28:70S–76S.
- Hinds D, Risch N. 1999. The ASPEX package: Affected sib-pair exclusion mapping.
- Hinrichs A, et al. 2005. Multipoint identity-by-descent computation for single-point polymorphism and microsatellite maps. *BMC Genet* 6(Suppl 1): S34.
- Hinrichs AL, et al. 2006. Functional variant in a bitter taste receptor (*hTAS2R16*) influences risk for alcohol dependence. *Am J Hum Genet* 78:103–111.
- Huang Q, Shete S, Amos CI. 2004. Ignoring linkage disequilibrium among tightly linked markers induces false-positive evidence of linkage for affected sib pair analysis. *Am J Hum Genet* 75:1106–1112.
- Iacono WG, et al. 1999. Behavioral disinhibition and the development of substance-use disorders: Findings from the Minnesota Twin Family Study. *Dev Psychopathol* 11:869–900.
- Iacono WG, et al. 2002. P3 event-related potential amplitude and the risk for disinhibitory disorders in adolescent boys. *Arch Gen Psychiatry* 59:750–757.
- Johnson EO, van den Bree MB, Pickens RW. 1996. Subtypes of alcohol-dependent men: A typology based on relative genetic and environmental loading. *Alcohol Clin Exp Res* 20:1472–1480.
- Kendler KS, et al. 2003. The structure of genetic and environmental risk factors for common psychiatric and substance use disorders in men and women. *Arch Gen Psychiatry* 60:929–937.
- Kendler KS, et al. 2006a. A joint genomewide linkage analysis of symptoms of alcohol dependence and conduct disorder. *Alcohol Clin Exp Res* 30:1972–1977.
- Kendler KS, et al. 2006b. A joint genomewide linkage analysis of symptoms of alcohol dependence and conduct disorder. *Alcohol Clin Exp Res* 30(12):1972–1977.
- Kessler RC, Borges G, Walters EE. 1999. Prevalence of and risk factors for lifetime suicide attempts in the national Comorbidity Study. *Arch Gen Psychiatry* 56:617–626.
- Krueger RF. 1999. The structure of common mental disorders. *Arch Gen Psychiatry* 56:1088–1096.
- Kuo PH, et al. 2006. Identification of susceptibility loci for alcohol-related traits in the Irish Affected Sib Pair Study of Alcohol Dependence. *Alcohol Clin Exp Res* 30(11):1807–1816.
- Lappalainen J, et al. 2005. Association between alcoholism and gamma-aminobutyric acid  $\alpha 2$  receptor subtype in a Russian population. *Alcohol Clin Exp Res* 29:493–498.
- Moskvina V, et al. 2009. Gene-wide analyses of genome-wide association data sets: Evidence for multiple common risk alleles for schizophrenia and bipolar disorder and for overlap in genetic risk. *Mol Psychiatry* 14(3):252–260.
- Moss HB, Lynch KG. 2001. Comorbid disruptive behavior disorder symptoms and their relationship to adolescent alcohol use disorders. *Drug Alcohol Depend* 64:75–83.
- O'hannessian CM, et al. 2004. The relationship between parental alcoholism and adolescent psychopathology: A systematic examination of parental comorbid psychopathology. *J Abnorm Child Psychol* 32:519–533.
- Pickens RW, et al. 1991. Heterogeneity in the inheritance of alcoholism. *Arch Gen Psychiatry* 48:19–28.
- Pickens RW, et al. 1995. Common genetic mechanisms in alcohol, drug, and mental disorder comorbidity. *Drug Alcohol Depend* 39: 129–138.
- Porjesz B, et al. 2005. The utility of neurophysiological markers in the study of alcoholism. *Clin Neurophysiol* 116:993–1018.
- Purcell S, et al. 2007. PLINK: A toolset for whole-genome association and population-based linkage analysis. *Am J Hum Genet* 81:559–575.
- Reich T. 1996. A genomic survey of alcohol dependence and related phenotypes: Results from the Collaborative Study on the Genetics of Alcoholism (COGA). *Alcohol Clin Exp Res* 20(8 Suppl): 133A–137A.
- Reich T, et al. 1998. Genome-wide search for genes affecting the risk for alcohol dependence. *Am J Med Genet* 81:207–215.
- Reynolds SM, et al. 2006. Neurotensin antagonist acutely and robustly attenuates locomotion that accompanies stimulation of a neurotensin containing pathway from rostral basal forebrain to the ventral tegmental area. *Eur J Neurosci* 24(1):188–196.

- Roberts SB, et al. 1999. Replication of linkage studies of complex traits: An examination of variation in location estimates. *Am J Hum Genet* 65:876–884.
- Slutske WS, et al. 1998. Common genetic risk factors for conduct disorder and alcohol dependence. *J Abnorm Psychol* 107:363–374.
- Slutske WS, et al. 2002. Personality and the risk for alcohol dependence. *J Abnorm Psychol* 111:124–133.
- Soyka M, et al. 2008. GABA-A2 receptor subunit gene (GABRA2) polymorphisms and risk for alcohol dependence. *J Psychiatr Res* 42(3): 184–191.
- Straub RE, et al. 1999. Susceptibility genes for nicotine dependence: A genome scan and followup in an independent sample suggest that regions on chromosomes 2, 4, 10, 16, 17 and 18 merit further study. *Mol Psychiatry* 4(2):129–144.
- Tarter RE, et al. 2004. Neurobehavior disinhibition in childhood predicts suicide potential and substance use disorder by young adulthood. *Drug Alcohol Depend* 76S:S45–S52.
- Wang JC, et al. 2004. Evidence of common and specific genetic effects: Association of the muscarinic acetylcholine receptor M2 (CHRM2) gene with alcohol dependence and major depressive syndrome. *Hum Mol Genet* 13:1903–1911.
- Wang S, et al. 2005. Whole-genome linkage analysis in mapping alcoholism genes using single-nucleotide polymorphisms and microsatellites. *BMC Genet* 6(Suppl 1): S28.
- Wang JC, et al. 2007. Functional variants in TAS2R38 and TAS2R16 influence alcohol consumption in high-risk families of African-American origin. *Alcohol Clin Exp Res* 31(2):209–215.
- Wetherill LF, et al. 2008. Neuropeptide Y receptor genes are associated with alcohol dependence, alcohol withdrawal phenotypes, and cocaine dependence. *Alcohol Clin Exp Res* 12:2031–2040.
- White HR, et al. 2001. Psychopathology as a predictor of adolescent drug use trajectories. *Psychol Addict Behav* 15:210–218.
- Wiener HW, et al. 2005. COGA phenotypes and linkages on chromosome 2. *BMC Genet* 6(Suppl 1): S125.
- Wilhelmsen KC, et al. 2005. Support for previously identified alcoholism susceptibility loci in a cohort selected for smoking behavior. *Alcohol Clin Exp Res* 29(12):2108–2115.
- Willour VL, et al. 2007. Attempted suicide in bipolar disorder pedigrees: Evidence for linkage to 2p12. *Biol Psychiatry* 61(5):725–727.
- Young SE, et al. 2000. Genetic and environmental influences on behavioral disinhibition. *Am J Med Genet* 96(5):684–695.
- Zubenko GS, et al. 2004. Genome-wide linkage survey for genetic loci that affect the risk of suicide attempts in families with recurrent, early-onset, major depression. *Am J Med Genet Part B* 129B(1): 47–54.