

### Diagnostic Testing Using Nasopharyngeal Swabs

#### INSTRUCTIONS ON NASOPHARYNGEAL SPECIMENS FROM HOSPITALIZED PATIENTS WITH SEVERE UNEXPLAINED FEBRILE RESPIRATORY ILLNESS

These laboratory instructions provide guidance on what specimens to collect and how to arrange transportation. It is absolutely critical that specimens be collected properly, labeled properly, stored refrigerated and submitted with a completed PHL laboratory submission form.

WHO: Hospitalized patients with severe unexplained febrile respiratory illness

**WHAT:** Preferred specimen is 1-2 nasopharyngeal swabs in 2-3 ml viral transport medium\* (VTM). Specimens should be collected within the first 24-72 hours of onset of symptoms, and no later than 5 days after onset of symptoms.

#### HOW:

#### Use infection control precautions:

- Personal protective equipment: fit-tested disposable N95 respirator [if unavailable, wear a medical (surgical mask)], disposable gloves, gown, and goggles.
- When completed, place all PPE in a biohazard bag for appropriate disposal.
- Wash hands thoroughly with soap and water or alcohol-based hand gel.

#### How to do a nasopharyngeal swab

- Use a flexible fine-shafted aluminum swab with a polyester (dacron or rayon, not cotton) tip.
- Insert swab into the nostril and back to the nasopharynx and leave in place for a few seconds.
- Slowly withdraw swab with a rotating motion.
- Place tip of the swab into a vial containing 2–3 ml of virus transport medium\* and cut the shaft.
- Label the transport tube with the patient first and last name, DOB, type of specimen, and <u>date and time</u> of collection.
- Complete the PHL virus isolation specimen submission form (http://www.ci.nyc.ny.us/html/doh/downloads/pdf/labs/labforms-virus\_id\_f2.pdf). \*\*Failure to complete or submit this form with the specimen could result in disposal of the specimen\*\*
- $\circ~$  A second swab should be used for the second nostril.
- $\circ$  Swabs of anterior nares and throat swabs are NOT acceptable for influenza testing.





#### **HOME ISOLATION**

#### Instructions for Suspected, Probable and Confirmed Swine Influenza Cases

Dear Patient,

You are suspected of having or were diagnosed recently with swine influenza (swine flu). Influenza can be transmitted when an infected person coughs or sneezes. It is also possible to become infected by touching surfaces contaminated with the virus, and then touching your own nose or mouth. Covering coughs and sneezes and washing hands with soap and water or with an alcohol-based hand rub are essential in stopping the spread of influenza.

During the period that you have symptoms, especially fever, it is possible for you to transmit this virus to those who have close contact with you (for example, someone living with or caring for you). To help prevent transmission of influenza virus to your close contacts, the Health Department is requesting that you **stay at home for 7 days after the day you developed a fever or 24-48 hours after all symptoms resolve, whichever is longer.** During that time, please follow these guidelines:

#### YOU SHOULD:

# Stay at home for 7 days after the first day of fever or at least 24-48 hours after all symptoms resolve, whichever is longer.

- Contact with others should be limited as much as possible; avoid close contact such as kissing, do not share toothbrushes, cigarettes or drinks with non-infected persons. Persons living in the home with you should limit contact with you to the extent possible. Consider designating one person (not someone who is pregnant) as the primary person who will care for you while you're sick.
- If possible, stay in a room separate from the common areas of the house. (For example, a spare bedroom, if that's possible.).
- ALWAYS cover your mouth and nose with a tissue when sneezing, coughing or blowing your nose and ALWAYS wash your hands with soap or water or use alcohol-based hand rubs after coughing or sneezing or throwing a used tissue in the garbage. Throw out your used tissues with your regular garbage. Do not touch your nose, face or eyes after coughing or sneezing.
- While the virus is likely not spread very well from contact with soiled household surfaces, if any of your body fluids (such as secretions from your nose or mouth, urine, vomit or stool) gets on surfaces in your home (such as door knobs or any other object that you sneeze or cough on), the surface should be washed with any standard household cleaner or disinfectant. Rinse the surface with tap water afterwards, and be sure to wash your hands after cleaning the area.
- Dirty dishes and eating utensils should be washed either in a dishwasher or by hand with warm water and soap. Separation of eating utensils for use by a patient with influenza is not necessary.
- Laundry can be washed in a standard washing machine with warm or cold water and detergent. It is not necessary to separate soiled linen and laundry used by a patient with influenza from other household laundry.

#### HOME ISOLATION - Instructions for Suspect, Probable and Confirmed Swine Influenza Cases (cont)

#### INSTRUCTIONS FOR CLOSE CONTACTS:

- Other household members should monitor themselves closely for the development of influenza-like symptoms. If they get sick with fever, cough, or sore throat, they should stay at home and take the same precautions listed above.
- Sick persons should not have visitors while they are ill with influenza or an influenza-like illness. If anyone who does not live with the patient must enter the home, they should avoid contact with the sick person.
- All persons in the household, especially those who must come in close contact with the sick person should clean their hands with soap and water or an alcohol-based hand rub frequently and after every contact with the sick person.
- If someone else is washing the used linens (such as bed sheets and towels) for the sick person, they should use household laundry soap and tumble dry on a hot setting. Avoid "hugging" laundry prior to washing it to prevent self-contamination and clean hands with soap and water or alcohol-based hand rub right after handling dirty laundry.
- Close contacts may consider using a facemask (e.g. surgical mask) or an N95 respirator. These can be purchased at a pharmacy or hardware store. If a reusable fabric facemask is used, it should be laundered with normal laundry detergent and tumble-dried in a hot dryer.

#### IF I AM SICK DO I NEED TO TAKE MEDICINE?

Right now the Health Department is not recommending that people with mild illness take any special medication for swine flu. You should rest, drink plenty of liquids, and take medications that you would normally take to treat your symptoms, such as acetaminophen or ibuprofen. People who are under 18 years of age should NOT take aspirin or aspirin-containing products (e.g., Pepto-Bismol<sup>™</sup>), as this can cause a rare serious illness called Reye's Syndrome in young people with the flu.

#### WHAT TO DO IF YOUR SYMPTOMS WORSEN

If your mild symptoms worsen, for example you develop difficulty breathing, you should seek medical attention from your own medical provider or at the emergency room. If you need to go to the doctor's office or emergency room, you should have a family member or friend drive you in a private car if at all possible, or take a taxi. Do NOT take public transportation (subway or bus). If you go to your doctor, please contact your doctor before you leave your home and tell the doctor you have influenza-like symptoms. You should go straight to the receptionist or triage nurse when you arrive and tell them that you have arrived so that you can be put in a private room while you wait to see the doctor. You may be asked to wear a mask. Try to sit away from others as much as possible. If you call an ambulance to take you to the hospital, let the operator know that you have influenza-like symptoms when you call 911, and let the ambulance crew know when they arrive.

### EMERGENCY WARNING SIGNS

Children: -Fast breathing or trouble breathing -Bluish skin color -Not drinking enough fluids Not waking up or not interacting -So irritable child doesn't want to be held -Flu-like symptoms improve but returns with fever and worse cough -Fever with a rash <u>Adults</u> -Difficulty breathing or shortness of breath Pain or pressure in the chest or abdomen -Sudden dizziness -Confusion -Severe or persistent vomiting

#### SHOULD PERSONS I LIVE WITH OR OTHER CLOSE CONTACTS TAKE MEDICINE TO PREVENT SWINE FLU?

The Health Department is recommending preventive medication particularly for household or other close contacts who have an underlying condition (such as certain chronic medical conditions, being immunocompromised, being <2 or > 65 years old) that may put them at risk for more severe illness. To find out whether preventive medication such as oseltamivir (Tamiflu<sup>®</sup>) is recommended, call a doctor.

#### WHERE TO GET MORE INFORMATION

For more information, call your doctor or visit the following websites that have information about swine flu: <a href="http://www.cdc.gov/www.who.int/www.nyc.gov/health">www.cdc.gov/www.who.int/www.nyc.gov/health</a> or call 311



Diagnostic Testing Using Nasopharyngeal Swabs (cont.)

#### Storage

- Place specimen(s) in a biohazard bag.
- Keep specimen(s) refrigerated at 4°C until pick-up.
- If specimens cannot be processed within 48-72 hrs, they should be kept frozen at or below –70 °C.

#### **Transportation to the Public Health Laboratory**

- o Insert the completed laboratory submission form.
  - Tips for completing the form:
    - Under DOHMH Bureau, check BCD
    - Under DOHMH Contact, put in the name of the contact person you spoke with
    - Under Test(s) requested, check Influenza A and B Direct Antigen Detection, Influenza A real-time RT-PCR, Influenza A Subtyping (H1/H3) real-time RT-PCR (see below)
    - Information on the form and the specimen has to match (name of pt, DOB, Specimen type, date and time of collection).
- Contact 1-866-NYC-DOH1 (Provider Access Line) to make a request to the DOHMH for pick-up.

4. TEST(S) REQUESTED*	
TX Influenza A and B Direct Antigen Detection	Rotavirus Direct Antigen Detection
Tall Influenza A and B real-time RT-PCR	RSV Direct Antigen Detection
Influenza A Subtyping (H1/H3) real-time RT-PCR	Virus Isolation/Detection
Influenza A/H5 real-time RT-PCR (DOHMH Request ONLY) <sup>1</sup>	Other (specify):
Smallpox Rule-Out Assays (DOHMH Request ONLY) <sup>1</sup>	Other (specify):
	LTR_DVI02_808

\* Virus transportation medium for use in collecting nasopharyngeal specimens

Add 10 g veal infusion broth and 2 g bovine albumin fraction V to sterile distilled water (to 400 ml). Add 0.8 ml gentamicin sulfate solution (50 mg/ml) and 3.2 ml amphotericin B (250 μg/ml). Sterilize by filtration.

THE CITY OF NEW YORK



DEPARTMENT OF HEALTH AND MENTAL HYGIENE Michael R Bloomberg Thomas R Frieden M.D. M.P.H.

Mauor

Thomas R. Frieden, M.D., M.P.H. Commissioner

nyc.gov/health

### 2009 New York City Department of Health and Mental Hygiene (NYC DOHMH) Health Alert #13: Swine Influenza Update

Please distribute to staff in the Departments of Critical Care, Emergency Medicine, Family Practice, Geriatrics, Internal Medicine, Infectious Disease, Infection Control, Pediatrics, Pharmacy, Neonatal Units, Pulmonary Medicine and Laboratory Medicine

#### April 28, 2009

PLEASE NOTE: This is a rapidly evolving situation. This alert provides interim guidance. Guidance is likely to change in the upcoming days and weeks as more information becomes available.

- Diagnostic testing at the Centers for Disease Control and Prevention (CDC) has now confirmed 44 cases of swine influenza A/H1N1 among students and staff at the St. Francis Preparatory High School in Queens (see case definitions below)
  - To date, all illness appears to be no more severe than usual seasonal influenza, and all patients we know of are recovering, with the overwhelming majority having had mild illness.
  - In some households with ill students, transmission to household members does appear to have occurred.
  - Treatment and prophylaxis recommendations are provided below.
- In addition, 5 new probable cases have been identified in New York City residents. Two of the probable cases involve people with links to Mexico, and three are from the St. Francis Preparatory High School community. Given the evolving epidemiology of this outbreak in NYC and elsewhere, the number of NYC cases is expected to increase in the coming days.
- Reporting and management of hospitalized patients in NYC with severe, unexplained febrile, respiratory illness:
  - Immediately report all patients with severe, unexplained febrile respiratory illness (e.g., pneumonia, acute respiratory distress syndrome, respiratory distress) to the Provider Access Line at 1-866-NYC-DOH1 (1-866-692-3641). Note that this is a change in the PAL number.
  - Test patients with severe febrile respiratory illness for influenza A using a commercially available rapid test, PCR or immunofluorescence test (e.g., DFA or IFA). If hospitals are not able to conduct initial rapid influenza testing, please contact the DOHMH PAL at 1-866-NYC-DOH1 (1-866-692-3641) to arrange for testing for influenza A. Note that the sensitivity of rapid tests may be less than 70%; patients who are rapid test negative may still have swine influenza.
  - In order to ensure sufficient laboratory resources are available to closely monitor for an increase in patients with severe, unexplained febrile respiratory illness in NYC, testing at the Public Health Laboratory will only be approved for cases that are first reported to DOHMH via the Provider Access Line. DOHMH staff will evaluate the case and advise whether testing for swine influenza at the Public Health Laboratory is indicated. DOHMH will facilitate specimen transport and testing for cases when testing is indicated.
- Management of patients with mild influenza-like illness in New York City
  - Patients with suspected (see epidemiologic risk factors listed below), probable or confirmed swine influenza and mild influenza-like illness (ILI) should stay home for 7 days after onset of symptoms, or until 24-48 hours after resolution of symptoms, whichever is longer.

Categories of urgency levels for NYC DOHMH Broadcast Notification System:

Health Alert: conveys the highest level of importance; warrants immediate action or attention Health Advisory: provides important information for a specific incident or situation; may not require immediate action Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action

- All patients with ILI, regardless of risk factors for swine influenza, should be instructed to stay home until at least 24-48 hours after their symptoms are resolved, wash their hands frequently, especially after coughing or sneezing, cough into a tissue (not into bare hands or onto another person), and dispose of tissues in the trash.
- At this time, we are <u>not</u> recommending influenza testing for persons with mild ILI.
- For patients with mild illness, treatment is only strongly recommended for people who also have underlying conditions that increase the risk for more severe illness due to influenza (listed below). For patients with mild illness who do not have underlying conditions, antiviral treatment can be offered but is NOT strongly recommended. In order to conserve supplies, providers are encouraged to be judicious in prescribing antivirals for uncomplicated mild ILI in patients without underlying conditions.
- Mild illness should only be treated if treatment can be started within 48 hours of symptom onset.
- Do <u>not</u> report cases of mild ILI to the DOHMH, unless they are associated with a cluster of illness (3 or more cases of ILI) in an institution such as a school, congregate living facility, or a long term care or medical facility.
- Infection control guidelines have been updated by CDC, and are available at <a href="http://www.cdc.gov/swineflu/guidelines\_infection\_control.htm">http://www.cdc.gov/swineflu/guidelines\_infection\_control.htm</a>
  - As always, medical facilities have screening and isolation protocols in place for patients presenting with fever and respiratory illness.
  - Signs should be posted asking all patients with fever and respiratory symptoms to notify medical staff immediately upon arrival.
  - Patients with influenza-like illness should be asked to don a face mask.
  - DOHMH requests that the following screening questions be asked at triage: During the 7 days prior to illness onset did you:
    - Travel to Mexico
    - Have close contact with an ill person associated with the St. Francis Preparatory High School in Queens, or
    - Have close contact with a known case of swine influenza.
  - At this time, patients who have ILI and any one of the above epidemiologic risk factors are considered suspected cases of swine influenza in New York City. This definition may be broadened at any time if wider community ILI transmission is documented.
  - Medical personnel caring for or obtaining respiratory specimens from patients with suspected, probable or confirmed swine influenza should take personal protective measures, including wearing a disposable fit-tested N95 mask and eye protection (goggles).
- Additional information on the outbreaks in the US and Mexico, including NYC, as well as further clinical guidance will be provided as it becomes available. For updated information on the national situation, see <a href="http://www.cdc.gov/swineflu/general\_info.htm">http://www.cdc.gov/swineflu/general\_info.htm</a>.

#### Dear Colleagues,

Testing at the CDC on April 26, 2009 has confirmed that an outbreak of influenza at the St. Francis Preparatory High School in Queens is due to swine influenza (H1N1). As of April 28, 2009, there are 44 confirmed cases among students and staff from the school. The DOHMH is actively investigating this outbreak and to date, all illnesses associated with the school appear to be mild. In addition, DOHMH has identified 5 probable cases of swine influenza, two of which are not associated with St. Francis Preparatory High School but are related to travel to Mexico.

As of April 28, 2009, the CDC has reported 64 laboratory confirmed human cases of swine influenza A/H1N1 in the US (45 in New York State, 10 in California, 6 in Texas, 2 in Kansas and 1 in Ohio). All 44 case patients in NYC have had illness that appears to be no more severe than seasonal influenza, with only one requiring brief hospitalization. All patients we know of in NYC appear to be recovering, and the overwhelming majority have had mild illness. No deaths have been reported. The virus is being described as

a new subtype of A/H1N1 not previously detected in swine or humans. Isolates from California and Texas have been found to be susceptible to the neuraminidase inhibitors (oseltamivir and zanamivir) but resistant to the adamantanes (amantadine and rimantadine). As of April 26, 2009, the Government of Mexico had reported 18 laboratory confirmed cases of swine influenza A/H1N1. Investigation is continuing to clarify the spread and severity of the disease in Mexico. Suspect clinical cases have been reported in 19 of the country's 32 states, including thousands of cases and more than 80 deaths – however many of these may be unconfirmed cases. Multiple other countries are also reporting cases. The World Health Organization yesterday raised the pandemic influenza alert level from phase 3 to phase 4.

The symptoms of swine influenza cases in the United States to date have been similar to routine seasonal influenza; they include fever, cough, sore throat, headache, chills, myalgias and fatigue. The incubation period is unknown at this time, but is likely similar to seasonal influenza (1-7 days). Anecdotally, the incubation period has been somewhat shorter, in the range of 1-3 days, but data are still being collected. The infectious period for a confirmed case of swine influenza A (H1N1) virus infection is defined as 1 day prior to the case's illness onset to 7 days after onset (day -1 to day 7). Viral shedding may be prolonged in children or the immunocompromised.

#### Surveillance for Swine Influenza in Hospitalized Cases Citywide

In order to determine whether the swine influenza virus is causing severe illness in New York City, DOHMH is focusing its surveillance efforts on <u>hospitalized patients with severe illness</u>.

DOHMH requests that providers immediately report any patient with severe unexplained febrile respiratory illness (e.g., pneumonia, ARDS, or respiratory distress) via the Provider Access Line 1-866-NYC-DOH1 (1-866-692-3641). These patients should be tested for influenza using either a commercial rapid test, or direct or indirect immunofluorescence. DOHMH will arrange for transportation of clinical specimens to the Public Health Laboratory for additional testing for swine influenza.

Note: The sensitivity of the rapid test for influenza is unknown and may be less than 70%. To reduce the likelihood that severe cases of swine influenza are missed, contact the health department to discuss testing of severely ill patients even if rapid influenza testing is negative.

**To allow DOHMH to prioritize laboratory testing, providers should not send specimens to the Public Health Laboratory unless advised to do so and arranged with DOHMH.** (See attached instructions for collecting and submitting laboratory diagnostic specimens for swine influenza testing.) Nasopharyngeal swabs are the preferred specimens for influenza testing in the current swine influenza context. Please note that strict personal protective measures should be taken when obtaining specimens, or providing patient care, including the donning of an disposable fit-tested N-95 mask, goggles, and placement of the patient in an airborne infection isolation room (AIIR), or if not available, a single room with a closed door.

#### What to report to the Health Department

Only patients with severe febrile respiratory illness of unknown etiology should be reported to the Health Department. Do not report patients with mild illness, even if there is a known risk factor for swine influenza.

#### Antiviral Treatment

## At this time antiviral treatment with oseltamivir or zanamavir is strongly recommended for the following:

- All hospitalized patients with suspected, probable or confirmed swine influenza (see case definitions below)
- All hospitalized patients with severe febrile unexplained respiratory illness (including ARDS, pneumonia or respiratory distress) pending testing for swine influenza.

- Patients with mild influenza-like illness AND underlying conditions (e.g., chronic pulmonary, cardiovascular, renal, hepatic, hematological or metabolic disorders, immunosuppression, compromised respiratory function, including conditions which increase the risk for aspiration, long-term aspirin therapy, pregnancy, age > 65 years, and age < 2 years) that increase the risk for more severe illness due to influenza. ]
- Treatment for any patient with mild ILI should only be started if within 48 hours of symptom onset.

## At this time, antiviral treatment with oseltamivir or zanamavir is NOT strongly recommended for the following:

- Patients with mild illness who <u>do not have</u> underlying conditions. For these individuals, antiviral treatment can be offered at the discretion of their provider. However, as above, treatment should only be started if within 48 hours of symptom onset.
- In order to conserve supplies and to minimize the emergence of resistance, providers should be judicious in prescribing antiviral medications to patients with mild ILI and no underlying conditions.

#### Antiviral Prophylaxis

Currently, DOHMH is recommending antiviral prophylaxis for the following persons:

- Healthcare workers who provided care to ill patients, and who either were not using or had a breach in appropriate personal protection when caring for or obtaining specimens from patients with influenza-like illness who have suspected, probable or confirmed swine influenza (see infection control guidance referenced below)
- Asymptomatic household and other close contacts of ill persons of suspected, probable or confirmed swine influenza cases who are either a) at higher risk for complications of influenza (listed below) or b) health care workers themselves.

#### Detailed guidance on antiviral therapy and prophylaxis is available at:

http://www.cdc.gov/swineflu/recommendations.htm, and at http://www.cdc.gov/flu/professionals/antivirals/dosagetable.htm#table

#### **Oseltamivir Supplies**

The possible increased demand for oseltamivir (Tamiflu®) may lead to shortages, and anecdotal reports suggest that the local supply may be limited. While there is a national and state stockpile of Tamiflu®, it is not readily available for use by the public. In order to avoid causing a shortage, we are *urging providers to use prudent judgment* when evaluating patients and deciding if antiviral treatment or prophylaxis is indicated.

#### Summary of Management of Persons with Mild Influenza-like Illness

At this time, providers assessing patients with mild ILI in clinical settings, including emergency departments, should not test for influenza. If these patients have already been diagnosed with probable or confirmed swine influenza, or if they have an epidemiologic risk factor for swine influenza, including travel to Mexico, close contact with a confirmed case of swine influenza or close contact with an ill person associated with St. Francis Preparatory High School, they should be sent home with instructions to stay at home for 7 days after onset of symptoms, or until 24-48 hours after their symptoms resolve, whichever is later, and instructed on the importance of hand and respiratory hygiene. Instructions should be given to seek medical care with worsening of symptoms (see signs of worsening illness below). Please see accompanying document regarding home isolation of patients with suspected swine influenza. This document should be provided to outpatients with suspected, probable or confirmed swine influenza upon discharge.

## Summary of Management of Persons with Mild Influenza-like Illness and Underlying Conditions that Increase the Risk of Severe Influenza Infection

Patients with mild ILI and underlying conditions placing them at higher risk for severe illness should be treated empirically for influenza. See <u>http://www.cdc.gov/swineflu/recommendations.htm</u> for specific guidelines. No specific testing for influenza is recommended. These patients may be sent home with instructions to stay at home for 7 days after onset of symptoms, or until 24-48 hours after their symptoms resolve, whichever is later, and instructed on the importance of hand and respiratory hygiene. Instructions should be given to seek medical care with worsening of symptoms.

These underlying conditions include chronic pulmonary, cardiovascular, renal, hepatic, hematological or metabolic disorders, immunosuppression, compromised respiratory function, including conditions which increase the risk for aspiration, long-term aspirin therapy, pregnancy, age  $\geq 65$  years, and age < 2 years.

#### **Updated Infection Control Recommendations**

Recommendations for infection control are evolving and have been revised almost daily. For detailed current infection control recommendations for medical facilities please see the attached Swine Influenza Q&A for Providers and the CDC website: <u>http://www.cdc.gov/swineflu/guidelines\_infection\_control.htm</u>.

Please note that for infection control purposes, NYC defines patients with suspected swine influenza as those who have acute febrile respiratory illness (regardless of severity), and an epidemiologic risk factor as described above. This differs from the current CDC suspected case definition. This definition may be broadened at any time if wider transmission is documented.

The Health Department requests that providers also immediately report any clusters of influenza-like illness (3 or more cases) in medical facilities, congregate settings such as long-term care facilities, or schools.

#### **Additional resources:**

\*CDC Swine Influenza Page - <u>http://www.cdc.gov/swineflu/</u> CDC Health Advisory - <u>http://www.cdc.gov/swineflu/pdf/HAN\_042509.pdf</u> NYC DOHMH Swine Flu Information - <u>http://www.nyc.gov/html/doh/html/cd/cd-swineflu.shtml</u> \*NYC DOHMH Home Page - <u>http://www.nyc.gov/html/doh/html/home/home.shtml</u> New York State Swine Flu Resources http://www.nyhealth.gov/diseases/communicable/influenza/seasonal/swine\_flu/index.htm

To contact the Health Department, including to report suspected cases of swine influenza in hospitalized patients and arrange for specimen testing, please call the Provider Access Line at 866-NYC-DOH1. This number is also available for questions or consultations by providers.

As always, we appreciate the cooperation of the medical community in New York City and will update you with further information when it becomes available.

Sincerely,

Annie Fine

Annie Fine, MD Medical Director Zoonotic, Influenza, & Vectorborne Diseases Unit Bureau of Communicable Disease

Scott Harper

Scott A. Harper, MD, MPH, MSc Medical Epidemiologist Zoonotic, Influenza, & Vectorborne Diseases Unit Bureau of Communicable Disease

#### New York City Case Definitions for Infection with Swine Influenza A (H1N1) Virus

- 1. <u>A Confirmed case of swine influenza A (H1N1) virus infection is defined as a person with influenza-like illness (ILI) with laboratory confirmed swine influenza A (H1N1) virus infection at CDC by one or more of the following tests:</u>
  - 1. real-time RT-PCR
  - 2. viral culture
  - 3. four-fold rise in swine influenza A (H1N1) virus specific neutralizing antibodies
- 2. <u>A Probable case</u> of swine influenza A (H1N1) virus infection is defined as a person with ILI with an influenza test that is positive for influenza A, but H1 and H3 negative.
- 3. <u>A Suspected case\*</u> of swine influenza A (H1N1) virus infection is defined as a <u>person with ILI who</u>, <u>during the 7 days prior to symptom onset</u>
  - 1. Traveled to Mexico
  - 2. Had close contact with an ill person associated with St. Francis Preparatory High School
  - 3. Had close contact with a confirmed or probable case of swine influenza.

\* Note that the NYC case definition for suspected swine influenza differs from the CDC case definition.

#### Conditions which increase the risk of severe influenza infection

- chronic pulmonary, cardiovascular, renal, hepatic, hematological, or metabolic disorders,
- immunosuppression,
- compromised respiratory function, including conditions which increase the risk for aspiration,
- long-term aspirin therapy
- pregnancy
- age <u>></u> 65 years
- age < 2 years

#### Signs and symptoms of worsening illness

In children emergency warning signs that need urgent medical attention include:

- Fast breathing or trouble breathing
- o Bluish skin color
- Not drinking enough fluids
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough
- Fever with a rash

In adults, emergency warning signs that need urgent medical attention include:

- Difficulty breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- $\circ$  Confusion
- Severe or persistent vomiting