



Contents lists available at SciVerse ScienceDirect

Addictive Behaviors



Genetic influences on craving for alcohol

Arpana Agrawal^{a,*}, Leah Wetherill^b, Kathleen K. Bucholz^a, John Kramer^c, Samuel Kuperman^d, Michael T. Lynskey^a, John I Nurnberger Jr.^b, Marc Schuckit^e, Jay A. Tischfield^f, Howard J. Edenberg^b, Tatiana Foroud^b, Laura J. Bierut^a

^a Washington University School of Medicine, USA^b Indiana University School of Medicine, USA^c University of Iowa School of Medicine, USA^d University of Iowa Hospitals and Clinics, USA^e University of California at San Diego, USA^f Rutgers University, USA

ARTICLE INFO

Available online xxx

Keywords:

Alcohol
Genes
Dopamine
GWAS
Genetics
Craving

ABSTRACT

Introduction: Craving is being considered for inclusion in the Diagnostic and Statistical Manual (DSM) DSM-5. However, little is known of its genetic underpinnings — specifically, whether genetic influences on craving are distinct from those influencing DSM-IV alcohol dependence.

Method: Analyses were conducted in a sample of unrelated adults ascertained for alcohol dependence (N = 3976). Factor analysis was performed to examine how alcohol craving loaded with the existing DSM-IV alcohol dependence criteria. For genetic analyses, we first examined whether genes in the dopamine pathway, including dopamine receptor genes (DRD1, DRD2, DRD3, DRD4) and the dopamine transporter gene (SLC6A3), which have been implicated in neurobiological studies of craving, as well as alpha-synuclein (SNCA), which has been previously found to be associated with craving, were associated with alcohol craving in this sample. Second, in an effort to identify novel genetic variants associated with craving, we conducted a genomewide association study (GWAS). For variants that were implicated in the primary analysis of craving, we conducted additional comparisons — to determine if these variants were uniquely associated with alcohol craving as compared with alcohol dependence. We contrasted our results to those obtained for DSM-IV alcohol dependence, and also compared alcohol dependent individuals without craving to non-dependent individuals who also did not crave alcohol.

Results: Twenty-one percent of the full sample reported craving alcohol. Of those reporting craving, 97.3% met criteria for DSM-IV alcohol dependence with 48% endorsing all 7 dependence criteria. Factor analysis found a high factor loading (0.89) for alcohol craving. When examining genes in the dopamine pathway, single nucleotide polymorphisms (SNPs) in DRD3 and SNCA were associated with craving ($p < 0.05$). There was evidence for association of these SNPs with DSM-IV alcohol dependence ($p < 0.05$) but less evidence for dependence without craving ($p > 0.05$), suggesting that the association was due in part to craving. In the GWAS, the greatest evidence of association with craving was for a SNP in the integrin alpha D (ITGAD) gene on chromosome 7 (rs2454908; $p = 1.8 \times 10^{-6}$). The corresponding p-value for this SNP with DSM-IV alcohol dependence was similar ($p = 4.0 \times 10^{-5}$) but was far less with dependence without craving ($p = 0.02$), again suggesting the association was due to alcohol craving. Adjusting for dependence severity (number of endorsed criteria) attenuated p-values but did not eliminate association.

Conclusions: Craving is frequently reported by those who report multiple other alcohol dependence symptoms. We found that genes providing evidence of association with craving were also associated with alcohol dependence; however, these same SNPs were not associated with alcohol dependence in the absence of alcohol craving. These results suggest that there may be unique genetic factors affecting craving among those with alcohol dependence.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

Genetic studies of alcohol dependence are at an exciting crossroad. With the emergence of the proposed DSM-5 criteria for diagnosing substance use disorders (dsm5.org), investigators with genetically

* Corresponding author at: Washington University School of Medicine, Dept. of Psychiatry, 660 S. Euclid, CB 8134, Saint Louis, MO 63110, USA. Tel.: +1 314 286 1778; fax: +1 314 286 2213.

E-mail address: arpana@wustl.edu (A. Agrawal).

informative data have the unique opportunity to revisit important hypotheses regarding the genetic underpinnings of specific components of alcohol use disorders.

Three important changes to the diagnosis of substance, including alcohol, use disorders are proposed in the DSM-5: (a) abuse and dependence may no longer be considered independent entities. Instead, a uni-dimensional score of 11 (abuse and dependence) criteria is proposed to diagnose moderate (2–3 criteria) and severe (4 or more criteria) substance use disorders; (b) “recurrent legal problems” may no longer be part of the criteria, due to its low frequency and weak correlation with the remaining abuse and dependence criteria; and (c) a new criterion, craving, has been suggested for addition to the existing criteria (O'Brien, 2011). While most investigators with data collected using DSM-IV criteria will experience limited problems with (a) and (b), the addition of a new criterion (c) can pose challenges for studies that have not collected data on craving. Particularly for large-scale genetic studies where it is often infeasible to conduct repeat data collection, the possibility exists that the addition of this new criterion will compromise genetic examination of the proposed DSM-5 defined substance use disorders. This report examines what additional genetic information craving contributes that cannot be gleaned from examining DSM-IV dependence criteria.

For DSM-5, craving is proposed to be defined as “strong desire or urge to use a specific substance”. The preliminary rationale for the addition of craving arose from three sources — its inclusion in the International Classification of Disease (ICD-10) (World Health Organization, 2007) definition of substance use disorders, neurobiological experiments supporting cue-elicited craving (Childress et al., 1999; Kalivas & O'Brien, 2008; Kalivas & Volkow, 2005; Volkow et al., 2006), and clinical observations of its significance in relapse (Flannery et al., 2001; Monti et al., 1990; O'Brien, 2005). There is little doubt that craving is an integral aspect of substance use disorders — numerous epidemiological (Keyes, Krueger, Grant, & Hasin, 2011; Mewton, Slade, McBride, Grove, & Teesson, 2011) and emergency room (Cherpitel et al., 2010) studies report high factor loadings for alcohol craving, indicating considerable correlation between self-reported craving and the underlying liability to substance use disorders as indexed by the DSM-IV dependence criteria. However, whether it adds unique information to the diagnosis of substance use disorder remains unclear.

In psychometric analyses, craving is considered a “difficult” item — that is, it is infrequently endorsed and even though it discriminates between individuals with and without substance use disorders, it does so primarily in individuals already at high liability to develop the disorder. In other words, those who report craving are typically already classified as substance dependent due to endorsement of multiple (often more than 3) dependence criteria; craving further discriminates these most severely affected individuals (i.e. those with multiple other symptoms) (Bohn, Krahm, & Staehler, 1995; Bucholz et al., 1996). Consequently, the addition of craving as a criterion does not produce a significant increase in those diagnosed with alcohol use disorders (Agrawal, Heath, & Lynskey, 2011; Cherpitel et al., 2010; Keyes et al., 2011; Mewton et al., 2011). This opens the door to the discussion of whether craving is best assessed using a single item or whether the architecture of craving is better suited to a multi-dimensional assessment in order to provide unique information beyond the existing definitions of dependence.

What does this mean for genetic studies of alcohol dependence? It is well established that genetic factors play a substantial role in the etiology of substance use disorders, regardless of the type of diagnostic classification used. Therefore, it is widely accepted that the proposed DSM-5 definitions of substance use disorder will also be heritable. However, the addition of this new item raises the question: are there novel genetic influences on craving? Even though phenotypic correlations between craving and other dependence criteria are high, this does not rule out the possibility that at least some genetic pathways that contribute to craving may be distinct from those influencing other aspects of substance dependence.

There are multiple routes for examining the potential role of distinct genetic influences on craving. In the literature, craving has been studied in the context of drug compulsion, and consequently as a correlate of relapse, particularly under negative emotional states (Marlatt & Gordon, 1985). Studies of cue-elicited craving report activation in brain regions that constitute dopamine pathways. For instance, Childress et al. (1999), reported increases in cerebral blood flow in limbic (amygdala and anterior cingulate) regions when cocaine users were exposed to drug videos. In addition, dopamine-rich regions such as the ventral tegmental area (VTA) have been implicated in the neuroplasticity underlying the development of addiction and the emergence of craving (Kalivas & O'Brien, 2008). Accordingly, genes in the dopamine pathway may yield promising insights regarding the genetic contributions to craving.

Based on this posited neurobiological basis of cue-elicited craving, we selected a set of genes that broadly constitute the dopamine receptor/transporter system. These included receptor-encoding genes (*DRD1*, *DRD2*, *DRD3*, *DRD4*) and the dopamine (*SLC6A3*, also known as *DAT*) transporter gene. We also selected α -synuclein (*SNCA*) which was previously implicated in a candidate gene association study of alcohol craving conducted on a partially overlapping sample (Foroud et al., 2007) as well as in an independent sample (Bonsch et al., 2005). While synucleins have been primarily studied in the context of Alzheimer's and Parkinson's disease (Irvine, El-Agnaf, Shankar, & Walsh, 2008), α -synucleins regulate dopamine D2 synthesis (Perez et al., 2002). Bonsch and colleagues found increased α -synuclein expression in alcoholics, which was positively correlated with alcohol craving scores (Bonsch et al., 2004; Bonsch et al., 2005). In addition to this focus on dopamine pathway genes, in an effort to identify novel genetic variants that were associated with alcohol craving, we conducted a genomewide association study (GWAS).

2. Methods

2.1. Sample

The sample used for this study was the Study of Addiction: Genes and Environment (SAGE: (Bierut et al., 2010)) which drew data from three independent family-based studies to create a sample of unrelated DSM-IV alcohol dependent cases and alcohol exposed controls. The parent studies were family-based and ascertained for alcohol dependence (Collaborative Study of the Genetics of Alcoholism, COGA, (Begleiter et al., 1995)), nicotine dependence (Collaborative Study of the Genetics of Nicotine Dependence, COGEND, (Bierut et al., 2007)) and cocaine dependence (Family Study of Cocaine Dependence, FSCD (Bierut, Strickland, Thompson, Afful, & Cottler, 2008)). The SAGE series consisted of 1899 unrelated DSM-IV alcohol dependent cases and 1938 unrelated alcohol exposed (drank at least one drink during the lifetime) controls.

2.2. Measures

All three parent studies used modified versions of the Semi-Structured Assessment for the Genetics of Alcoholism (SSAGA, Bucholz et al., 1994) to assess substance use disorders with good validity (Hesselbrock, Easton, Bucholz, Schuckit, & Hesselbrock, 1999) and reliability (Bucholz et al., 1994). Alcohol craving was evaluated with a single item. In COGA and COGEND, the item queried “In situations where you could not drink alcohol, did you ever have such a strong desire for it that you couldn't think of anything else?” In FSCD, the item queried “Have you ever had such a strong craving for alcohol that you couldn't think of anything else?” DSM-IV alcohol dependence was defined by the clustering of 3 or more dependence criteria in a 12 month period. Note that, consistent with DSM-IV, craving was not utilized as a criterion to diagnose DSM-IV alcohol dependence.

2.3. Genotypic data

Genotyping was done with the Illumina Human 1 M beadchip by the Center for Inherited Diseases Research (CIDR) at Johns Hopkins University. A thorough data cleaning procedure on the genotype samples was applied to ensure the highest possible data quality, including using HapMap controls, detection of gender and chromosomal anomalies, hidden relatedness, population structure, missing call rates, plate effects, duplication error detection, and Hardy–Weinberg equilibrium (excluding SNPs with $p < 10^{-6}$) (Bierut et al., 2010; Laurie et al., 2010). The final number of autosomal and X-chromosome markers consisted of 948,142 SNPs (only autosomal SNPs were utilized). The software package EIGENSTRAT/EIGENSOFT (Price et al., 2006) was used to calculate principal components reflecting continuous variation in allele frequencies representing ancestral differences in subjects. Two principal components distinguishing African-American participants from European-American participants and Hispanic and non-Hispanic subjects, were identified (Bierut et al., 2010).

2.4. Factor analyses

Descriptive statistics were computed in SASv9 (SAS Institute, 1999). As the items assessing craving in COGA/COGEND and FSCD were slightly different, a multi-group confirmatory factor model was used to determine (a) the factor loading (discrimination) and threshold (difficulty) for craving in the full SAGE sample and (b) to examine whether these parameters varied across the 3 contributing parent studies. Item response modeling was conducted in MPlus v5 (Muthen & Muthen, 2007).

2.5. Statistical analysis

In the primary analysis, individuals endorsing craving (with or without DSM-IV alcohol dependence) were designated as affected and those who did not report craving (whether or not they met criteria for DSM-IV alcohol dependence) were designated as unaffected. Secondary analyses were performed to examine the association of particular SNPs with alcohol dependence without craving. For this analysis, individuals meeting criteria for DSM-IV alcohol dependence who did not endorse the proposed DSM-5 craving item were designated as affected and those who did not report craving and did not meet criteria for DSM-IV dependence were designated as unaffected. Results were also compared to those from our previous GWAS for DSM-IV alcohol dependence (Bierut et al., 2010).

PLINK (Purcell et al., 2007) was used to conduct all autosomal association analyses using logistic regression models. Genotype was coded log-additively (AA, Aa/aA, aa) and covariates representing sex, age at interview, whether participants were drawn from COGA or FSCD (COGEND serving as the reference group) and the first two principal components to account for ethnic variation in allele frequencies, were also included. As our primary analysis, we examined SNPs in *SNCA*, which has been previously studied in relation to alcohol craving, as well as genes in the dopamine pathway (*DRD1*, *DRD2*, *DRD3*, *DRD4*, *SLC6A3*). For the analysis of these genes of *a priori* significance, we set a liberal threshold of $p < 0.05$. Next, we conducted a genome-wide search — as these analyses are exploratory, statistical significance in the GWAS was set at $p < 5 \times 10^{-8}$. For both association analyses, primary analyses were conducted to identify SNPs associated with alcohol craving. For SNPs that were implicated in the primary analysis of craving, secondary analyses were performed examining other comparisons. These included an analysis of only non-craving individuals, contrasting those with and without alcohol dependence, as well as an analysis of DSM-IV alcohol dependence and an adjustment for severity of alcohol dependence.

3. Results

3.1. Sample characteristics

Table 1 provides the characteristics of those endorsing craving compared with those who did not endorse it. An overwhelming proportion (97.3%) of those endorsing craving met criteria for DSM-IV dependence. Nearly 48% of those who reported alcohol craving also endorsed all 7 DSM-IV dependence criteria. COGA contributed a majority of the craving cases for alcohol. Alternatively, as shown in Fig. 1, endorsement of craving increased with increasing number of dependence criteria endorsed with nearly 80% of those who reported all 7 DSM-IV criteria also reporting craving.

3.2. Factor analysis

Standardized factor loadings for all dependence criteria (0.80–0.92) and for craving (0.89) were high in the full sample (Table 2). When factor loadings and thresholds (estimates of prevalence) for craving were allowed to vary across COGA, COGEND and FSCD, factor loadings remained consistently high while thresholds varied modestly across samples. A 4 degree of freedom test resulted in a significant chi-square difference ($\Delta\chi^2 = 27.8$) although this difference was entirely attributed to the threshold differences. Furthermore, excluding craving from the factor analysis did not perturb the factor loadings for the DSM-IV dependence criteria.

3.3. Genes in the dopamine pathway

Table 3 shows the results for SNPs in genes in the dopamine pathway. Only SNPs with a p-value less than 0.05 for craving are shown and only for this subset of SNPs were comparisons with DSM-IV alcohol dependence and dependence without craving made. Setting a liberal threshold of $p < 0.05$, SNPs in *DRD2*, *DRD3* and *SNCA* showed evidence for association with craving. Comparing the results for these SNPs with those for DSM-IV alcohol dependence, SNPs in *DRD3* and *SNCA* appeared to have independent genetic effects on craving, infrequently showing a p-value < 0.05 for alcohol dependence while for *DRD2*, a number of SNPs associated with craving had a p-value < 0.05 for alcohol dependence. With some exception, the SNPs associated with craving did not show p-values < 0.05 for dependence without craving.

3.4. GWAS

Fig. 2 and Table 4 summarize the results of the autosomal genome-wide analysis using the craving for alcohol phenotype. The best association for craving was found for SNP rs2454908 (p-value = 1.83×10^{-6})

Table 1

Sample characteristics for 3976 individuals from the Study of Addiction: Genes and Environment.

	Craving (N = 841)	No craving (N = 3135)	
		Alcohol dependence (N = 1080)	No alcohol dependence (N = 2055)
Sex (male)	59.2%	61.8%	31.2%
Mean age	39.4 [SD 9.2]	38.6 [SD 9.3]	39.3 [SD 9.1]
European-American	67.5%	62.9%	74.0%
Hispanic	4.4%	3.6%	3.1%
COGA	57.1%	38.8%	26.6%
FSCD	25.1%	31.9%	29.8%
COGEND	17.8%	29.4%	45.2%
Meet criteria for DSM-IV dependence	97.3%	100%	0%
Mean number of dependence criteria	5.95 [SD 1.3]	4.5 [SD 1.3]	0.6 [SD 0.8]

SD — standard deviation.




Fig. 1. Percentage (dark shading) of those with 0 to 7 DSM-IV dependence criteria who also endorsed alcohol craving.

Table 2
Factor loadings and endorsement prevalence for craving relative to DSM-IV dependence criteria in 3976 subjects from SAGE.

	Loading	Prevalence (%)
Craving	.89	21
DSM-IV criteria		
Tolerance	.80	47
Withdrawal	.91	25
Larger/longer	.87	61
Quit attempts	.87	46
Time spent	.89	28
Give up activities	.92	29
Use despite problems	.88	46

The factor loadings are drawn from a confirmatory single factor model fit in MPlus.

on chromosome 16. Table 4 also compares results for craving with those for DSM-IV alcohol dependence and with dependence without craving. For rs2454908, the odds-ratio and p-value for craving and DSM-IV dependence were comparable; however, the p-value for this SNP was 0.0186 when examining dependence without craving. Of the 30 most significant autosomal SNPs for craving, 20 SNPs also had p-values less

than 0.05 for DSM-IV alcohol dependence with or without craving, but only rs2454908 had a p-value less than 0.05 for dependence without craving. The overlap between alcohol dependence and craving dropped steadily as the threshold for association with craving was made less stringent (Fig. 3).

3.5. Adjusting for severity of alcohol dependence

As indicated in Fig. 1, craving was more common in those endorsing a greater number of alcohol dependence criteria, an index of severity. We re-ran association analyses for SNPs in Tables 3 and 4 including number of dependence criteria (0–7) as a covariate. Adjustment for symptom count, as expected, attenuated p-values, in some cases. For genes in the dopamine pathway, after correction, the 3 *DRD3* SNPs with the lowest p-values (rs167770, rs226082 and rs324029) continued to show p-values less than 0.05. Three SNPs in *SNCA* also remained associated (rs6834765, rs11944331 and rs3775439) with p-values indicating a greater degree of significance upon adjustment for severity. For the GWAS, only 2 SNPs (rs1595411 and rs10115971) were no longer associated with craving at $p \leq 0.05$ after adjusting for severity. For example, after accounting for number of alcohol dependence criteria, rs2454908 retained a p-value of 0.0002.

4. Discussion

Craving, reflecting strong desire or urge to use a drug, is being proposed for inclusion in the DSM-5. In the present analyses, while we did not identify any genome-wide association signals for craving, we found that a number of SNPs that were associated with alcohol craving at p-values less than 1×10^{-6} also showed nominal association with DSM-IV alcohol dependence, but not with dependence excluding craving. Several SNPs in dopamine-related genes did exhibit differing results when comparing craving and DSM-IV alcohol dependence albeit at far lower levels of statistical significance. While imposing a high statistical threshold yielded significant overlap across SNPs for craving and for alcohol dependence, lowering this threshold and specifically focusing on

Table 3
Genes in the dopamine receptor/transporter pathway (only SNPs at $p \leq 0.05$ for craving are shown).

SNP		Craving (with or without alcohol dependence).				DSM-IV dependence		Dependence without craving	
	Position (base pair)	O.R.	Lower C.I.	Upper C.I.	p-Value	O.R.	p-Value	O.R.	p-Value
DRD3, chromosome 3, 27 SNPs									
rs9828046	115340953	0.71	0.53	0.94	1.65E-02	0.78	2.93E-02	0.93	5.71E-01
rs2630349	115356062	1.19	1.00	1.40	4.61E-02	1.01	9.04E-01	0.97	7.23E-01
rs167770	115362252	0.82	0.73	0.93	2.07E-03	0.97	6.15E-01	1.09	1.79E-01
rs226082	115363703	0.82	0.72	0.93	1.82E-03	0.98	7.65E-01	1.10	1.26E-01
rs324029	115364313	0.82	0.73	0.93	1.93E-03	0.98	6.80E-01	1.09	1.66E-01
SNCA, chromosome 4, 31 SNPs									
rs35101723	90866209	0.32	0.11	0.91	3.23E-02	0.41	1.09E-02	0.53	9.00E-02
rs6834765	90903013	0.83	0.69	0.98	3.14E-02	0.96	5.51E-01	1.05	6.07E-01
rs3796661	90906530	0.76	0.58	0.98	3.63E-02	0.83	8.66E-02	0.92	4.99E-01
rs11944331	90909352	0.84	0.71	0.99	3.23E-02	0.98	7.38E-01	1.09	3.10E-01
rs3775439	90928764	0.86	0.74	0.99	3.28E-02	0.93	2.68E-01	1.02	7.55E-01
DRD2, chromosome 11, 34 SNPs									
rs1076560	112788898	1.20	1.02	1.41	2.72E-02	1.13	8.72E-02	1.08	3.71E-01
rs2734838	112791711	0.88	0.78	0.99	3.55E-02	0.87	8.60E-03	0.88	3.92E-02
rs1079727	112794392	1.21	1.03	1.42	2.22E-02	1.13	1.04E-01	1.06	4.74E-01
rs4986918	112800452	0.55	0.34	0.89	1.48E-02	1.08	6.89E-01	1.30	1.81E-01
rs1076563	112801119	0.87	0.77	0.98	2.67E-02	0.87	7.15E-03	0.88	3.68E-02
rs1116313	112801317	0.87	0.77	0.98	2.04E-02	0.86	6.01E-03	0.88	3.46E-02
rs12364051	112810524	0.88	0.78	1.00	4.29E-02	0.88	2.14E-02	0.89	6.90E-02
rs17529477	112822277	0.84	0.73	0.96	1.31E-02	0.91	9.24E-02	0.97	6.21E-01

O.R. odds-ratio; lower/upper C.I. – lower and upper 95% confidence interval on odds-ratio. Position in base pair and gene based on NCBI build 36.3.




Fig. 2. Manhattan plot showing genomewide association study (GWAS) results for alcohol craving.

genes of *a priori* interest indicated potential biological specificity for craving.

We selected genes in the dopamine receptor/transporter pathways because multiple imaging studies have implicated dopamine-rich brain regions in experiments evoking cue-elicited craving for cocaine (Childress et al., 1999; Volkow et al., 2006) and other drugs (Kalivas & O'Brien, 2008; Kalivas & Volkow, 2005). This limbic activation suggests that genes in dopamine pathways might be of salience to craving but not all aspects of alcohol dependence. Despite the significant overlap between DSM-IV alcohol dependence and craving in our sample, we saw some support for differential effects of SNPs on craving, alcohol dependence and dependence without craving. We also found support for

signals in SNCA, a gene previously implicated in a partially overlapping sample (Foroud et al., 2007). However, the SNPs identified from this study were not the same as those identified by Foroud and colleagues.

Multiple SNPs in *DRD3* were associated with craving alone, which supports the role of dopamine pathway genes in craving. Dopamine receptors in the striatum are down-regulated in alcohol dependent individuals (Volkow et al., 1996), and this reduction in striatal D2/3 receptors was shown to be associated with increased craving in alcoholics (Heinz et al., 2004). These authors subsequently demonstrated that the increased alcohol craving was inversely related to dopamine synthesis and binding in the same subjects (Heinz et al., 2005). Similarly, Tupala and colleagues reported reduced D2/D3 binding in the nucleus accumbens and amygdala among late-onset type 1 alcoholics, who have been hypothesized to have a deficit in dopamine availability (Tupala et al., 2001). A more general study revealed that decreased striatal D2/3 availability in healthy controls was associated with greater impulsivity and craving for stimulants (Buckholtz et al., 2010). D3 knock-out mice also show intense withdrawal from ethanol (Narita, Soma, Tamaki, Narita, & Suzuki, 2002). In fact dopamine activation has been proposed to treat addictions in general, (Blum, Liu, Shriner, & Gold, 2011) and more specifically, as a therapeutic target to reduce craving and relapse in the “dopamine-impooverished addicted brain” (Diana, 2011). Our results, despite not achieving genomewide levels of statistical significance, are promising in that they support the results reported above.

In reviewing the present findings, it should also be recognized that craving experienced as a consequence of cue exposure may differ from self-reported “strong desire or urge” to use alcohol, which is the phenotype used in this study (Carter & Tiffany, 1999; Kozlowski, Mann, Wilkinson, & Poulos, 1989; Marlatt & Gordon, 1985). Furthermore, craving is commonly reported during abstinence and can be an aspect of

Table 4

Top 30 signals from GWAS of alcohol craving with corresponding results for the top SNPs from GWAS of DSM-IV alcohol dependence and dependence without craving.

Chromosome	SNP	Position (base pair)	Gene	Craving				DSM-IV dependence		Dependence without craving	
				O.R.	Lower C.I.	Upper C.I.	p-Value	O.R.	p-Value	O.R.	p-Value
16	rs2454908	31318686	<i>ITGAD</i>	0.74	0.65	0.83	1.83E-06	0.80	4.04E-05	0.86	1.86E-02
14	rs11849247	87778256	<i>KCNK10</i>	1.67	1.34	2.08	4.44E-06	1.31	1.05E-02	1.15	2.70E-01
9	rs3849883	30629205		2.41	1.65	3.51	5.45E-06	1.51	3.91E-02	1.05	8.51E-01
8	rs2469518	133225649	<i>KCNQ3</i>	0.77	0.68	0.86	7.06E-06	0.87	8.31E-03	0.96	4.95E-01
7	rs6466086	105550662		1.41	1.21	1.65	1.26E-05	1.17	2.47E-02	1.04	6.10E-01
17	rs34870220	7025635		1.81	1.39	2.36	1.26E-05	1.34	2.20E-02	1.00	9.99E-01
8	rs1991718	133230506	<i>KCNQ3</i>	0.77	0.69	0.87	1.60E-05	0.85	1.33E-03	0.92	1.71E-01
11	rs2447947	75621188		1.29	1.15	1.44	1.60E-05	1.13	1.48E-02	1.02	7.51E-01
3	rs4561856	87742811		0.73	0.63	0.84	1.79E-05	0.91	1.44E-01	1.04	5.53E-01
19	rs198967	56104833	<i>KLK4</i>	2.02	1.47	2.79	1.84E-05	1.17	3.30E-01	0.83	3.31E-01
6	rs7762384	145081988	<i>UTRN</i>	0.59	0.46	0.75	2.03E-05	0.98	8.35E-01	1.24	6.15E-02
16	rs17855121	30911670	<i>STX1B2</i>	1.32	1.16	1.50	2.21E-05	1.17	6.79E-03	1.06	4.15E-01
12	rs721793	66720281		1.40	1.20	1.64	2.28E-05	1.25	2.50E-03	1.07	4.40E-01
16	rs767505	55770151	<i>NIP30</i>	0.70	0.60	0.83	2.36E-05	0.78	3.10E-04	0.87	7.39E-02
3	rs11707897	2589626	<i>CNTN4</i>	0.68	0.56	0.81	2.41E-05	0.82	8.00E-03	0.93	4.02E-01
14	rs4983262	27190834		0.75	0.65	0.86	2.44E-05	0.88	3.02E-02	0.98	8.11E-01
19	rs198966	56103822	<i>KLK4</i>	1.85	1.39	2.46	2.61E-05	1.17	2.73E-01	0.87	4.17E-01
2	rs1901439	134153941		0.69	0.58	0.82	2.67E-05	0.87	5.15E-02	1.05	5.68E-01
7	rs17132242	2145988	<i>MAD1L1</i>	1.50	1.24	1.81	2.80E-05	1.16	9.93E-02	0.96	7.04E-01
11	rs10899187	75643975		1.31	1.15	1.48	2.81E-05	1.17	6.56E-03	1.06	3.97E-01
12	rs3782323	48769188	<i>SMARCD1</i>	0.76	0.67	0.86	2.83E-05	0.90	6.65E-02	1.02	7.54E-01
4	rs2666055	179581414		1.44	1.22	1.72	2.90E-05	1.16	6.63E-02	1.03	7.81E-01
8	rs1595411	133228607	<i>KCNQ3</i>	0.78	0.70	0.88	3.12E-05	0.85	1.17E-03	0.92	1.37E-01
2	rs687415	166011844	<i>TTC21B</i>	1.41	1.20	1.65	3.15E-05	1.19	2.53E-02	1.06	5.34E-01
20	rs8114108	24980075	<i>ACSS1</i>	0.59	0.46	0.76	3.15E-05	0.92	4.53E-01	1.11	3.82E-01
17	rs12947697	7030341		1.76	1.35	2.30	3.19E-05	1.30	4.17E-02	0.96	8.12E-01
9	rs10115971	100064708		1.35	1.17	1.56	3.37E-05	1.18	1.16E-02	1.07	4.08E-01
16	rs17790906	30477160	<i>MGC13138</i>	1.50	1.24	1.82	3.73E-05	1.24	1.80E-02	1.05	6.48E-01
6	rs6899567	157595		0.69	0.58	0.82	3.76E-05	0.93	3.35E-01	1.07	4.53E-01
17	rs12603684	74735394		1.33	1.16	1.53	4.02E-05	1.16	1.88E-02	1.03	6.45E-01

O.R. odds-ratio; lower/upper C.I. – lower and upper 95% confidence interval on odds-ratio. Position in base pair and gene based on NCBI build 36.3.




Fig. 3. Proportion of SNPs with a p -value ≤ 0.05 for DSM-IV alcohol dependence among SNPs with varying p -values for alcohol craving.

withdrawal, but whether anticipatory pleasure from alcohol use, cue-induced urges, desires to drink during withdrawal and self-reported craving are overlapping, or even related, constructs has not been well-explored (Moss, 2011). There is also evidence that craving for some drugs (e.g. cocaine) is a significant contributor to relapse (O'Brien, 2005) with one study reporting that alcohol craving was associated with time to relapse after residential treatment (Oslin, Cary, Slaymaker, Collieran, & Blow, 2009). However, this effect was not noted for alcohol in one large study (Leggio, Ray, Kenna, & Swift, 2009) and the effects were found to be even weaker when considering self-reported craving (Drummond, 2001; Drummond, Litten, Lowman, & Hunt, 2000; Lowman, Hunt, Litten, & Drummond, 2000).

It should also be noted that cue-elicited craving may be accentuated in individuals experiencing negative emotional states, particularly in high risk situations. In the context of the cognitive behavioral relapse prevention model (Marlatt & Gordon, 1985), under situations of stress (Hendershot, Witkiewitz, George, & Marlatt, 2011; Larimer, Palmer, & Marlatt, 1999), the anticipatory pleasure (i.e. positive reinforcement, preexisting expectancies) associated with drug-taking is heightened, intensifying craving. In fact, stress-induced craving has been found to be a more potent predictor of relapse than drug-cue induced craving (Sinha, Garcia, Paliwal, Kreek, & Rounsaville, 2006; Sinha et al., 2011). Therefore, our association signals in dopamine pathway genes, which have previously been implicated in cue-induced craving, may have shown modest p -values for self-reported craving, in part, due to the well-documented variations between cue-induced, stress-induced and self-reported craving.

The most significant association for alcohol craving in this GWAS was an intronic SNP, rs2454908, in the integrin, alpha D (*ITGAD*) gene which belongs to the CD11 antigen-like family of genes. To the best of our knowledge, the only potentially functionally relevant SNP that is in moderate linkage disequilibrium with this marker is in the 3' untranslated regulatory region of the neighboring Integrin, alpha X (*ITGAX*) gene. How this SNP relates to alcohol craving is unknown. A number of SNPs in the potassium voltage-gated channel, KQT-like subfamily, member 3 (*KCNQ3*) gene were also associated with craving. The protein encoded by this gene, along with the protein product of *KCNQ2* and *KCNQ5*, forms potassium M channels that are critical in neuronal excitability and variants in *KCNQ5* were previously identified in an African-American subset of individuals with alcohol dependence (Kendler et al., 2011). Of particular relevance to our study, Koyama, Brodie, and Appel (2007) have reported that ethanol induced excitation of dopamine neurons in the Ventral Tegmental Area (VTA) can be attributed to inhibition of currents in M-channels.

An important observation in these data was that (a) nearly all those who endorsed craving met criteria for DSM-IV alcohol dependence and (b) craving was more commonly reported in those with a greater

number of alcohol dependence symptoms, which is consistent with a prior latent class analysis in an extended subset of these data (Foroud et al., 1998). These findings, despite being drawn from a study ascertained for alcoholism, are consistent with the extant epidemiological literature that notes that craving, while being an excellent item when distinguishing between those at high versus lower liability to alcohol dependence, is only effective at making this distinction in individuals at high risk. In the context of psychometrics, the craving item has high discrimination (i.e. good factor loading) and high difficulty/severity (i.e. infrequently endorsed by low risk individuals). For instance, in the National Longitudinal Alcohol Epidemiologic Survey (NLAES), craving cohered well with other DSM-IV alcohol dependence criteria, had the second highest discrimination and also the highest severity (with the exception of the abuse criterion of legal problems, which is proposed for exclusion in the DSM-5). This indicates that the utility of craving will be most noticeable in distinguishing individuals with multi-symptom alcohol dependence but less so in those at low to moderate risk. That craving will add additional alcohol use disorder cases to DSM-5 diagnosis remains uncertain as, in our study and other epidemiological investigations (e.g. (Agrawal et al., 2011; Keyes, Krueger, Grant, & Hasin, 2010)), those reporting craving already met the burden of diagnosis. Therefore, as currently conceptualized craving, doesn't significantly modify the epidemiology of alcohol use disorders but adds predictive utility in high risk individuals. Despite this significant overlap between craving and severity of alcohol dependence, a number of our candidate gene and a majority of the GWAS signals retained p -values ≤ 0.05 after adjusting for severity. This would suggest further specificity of these SNPs in the etiology of craving.

Some limitations of our study should be noted. First, our sample drew from parent studies with differing ascertainment criteria and this may have contributed to heterogeneity in those endorsing craving. Our factor analyses, however, indicated that heterogeneity is attributable to the relative frequency of endorsement of craving across the parent studies, but not in its relationship with dependence. Second, as our sample was designed to be enriched for alcohol and drug dependent individuals, results from our study may not generalize to the population. Nonetheless, phenotypic similarities with epidemiological studies are apparent. Third, none of the association signals surpassed genomewide significance and hence, our comparison across craving, dependence without craving and DSM-IV dependence, which relied on p -values, may be confounded with a high rate of false positive findings. Replication will be necessary before we can be confident in these findings. Although 97% of individuals endorsing the craving item were alcohol dependent, the observation that some SNPs uniquely associated with craving while other SNPs were only associated with alcohol dependence demonstrates both the heterogeneity of the alcohol dependence diagnosis and the potential specificity of the craving item. Similar results were reported by Foroud et al. (2007) in that SNPs in *SNCA* were associated with craving but not with alcohol dependence.

Perhaps most importantly, the definition of craving is known to have a significant impact on its psychometric relationship with the construct of dependence — we used a single item that approximates the proposed DSM-5 criteria, however more work is required to outline whether this definition of craving is optimal for diagnostic purposes and for genetic analysis. The single craving item was employed in both a family-based study of European subjects in which it was associated with SNPs in *SNCA* (Foroud et al., 2007) as well as in a family-based linkage study of Mission Indians, which estimated the heritability of craving using this single item to be 65% (Ehlers & Wilhelmsen, 2005). Although a multi-dimensional craving assessment such as the Desires for Alcohol Questionnaire (DAQ) might offer a more powerful model providing overlapping and perhaps additional independent association results, we were unable to employ this measure in the current study because the questionnaire was only administered to a small subset of one participating study. However, Kramer et al. (2010) have studied the DAQ and found it to correlate well with this single craving item. Whether the 3

facets identified in that study (Strong Desires/Intentions, Negative Reinforcement and Positive Reinforcement and ability to control drinking) yield differing genetic signals should be investigated when larger samples with genetic data become available.

Overall, in our sample ascertained for alcohol dependence, we found that craving was strongly associated with severity of dependence. This may indicate its potential role in relapse. However, based on our work and that of numerous others, there is a significant need for research to further clarify how craving should be operationalized for diagnostic and research purposes. The genomic analyses presented here lead to two important conclusions – first, much larger sample sizes will be required to detect genes of small effect for craving. Despite the relatively high prevalence for craving in our sample (relative to epidemiological cohorts), we did not succeed in identifying signals of genomewide significance. Second, despite the overlap in some of the top signals across craving and alcohol dependence, there was some evidence of specificity in association when examining genes in the dopamine pathway and further, our GWAS signals for craving do not appear to be entirely attributable to the high degree of overlap between craving and severity of alcohol dependence. There needs to be a continued effort to examine candidate genes that may be particularly relevant to craving – while we explore one major family of genes in this study, multiple others in GABA-ergic, glutamatergic and opioidergic pathways have also been posited as participating in the neurobiology of craving (Kalivas & Volkow, 2005). While we could not conclude the extent to which the specificity of our association signals for the dopamine genes was due to false positive or negative findings, future studies may wish to focus on these genes with additional measures of craving.

Role of funding sources

Funding support for the Study of Addiction: Genetics and Environment (SAGE) was provided through the NIH Genes, Environment and Health Initiative [GEI] (U01 HG004422). SAGE is one of the genome-wide association studies funded as part of the Gene Environment Association Studies (GENEVA) under GEI. Assistance with phenotype harmonization and genotype cleaning, as well as with general study coordination, was provided by the GENEVA Coordinating Center (U01 HG004446). Assistance with data cleaning was provided by the National Center for Biotechnology Information. Support for collection of datasets and samples was provided by the Collaborative Study on the Genetics of Alcoholism (COGA; U10 AA008401), the Collaborative Genetic Study of Nicotine Dependence (COGEND; P01 CA089392), and the Family Study of Cocaine Dependence (FSCD; R01 DA013423). Funding support for genotyping, which was performed at the Johns Hopkins University Center for Inherited Disease Research, was provided by the NIH GEI (U01HG004438), the National Institute on Alcohol Abuse and Alcoholism, the National Institute on Drug Abuse, and the NIH contract “High throughput genotyping for studying the genetic contributions to human disease” (HHSN268200782096C). Other support includes DA23668 and ABMRF/Foundation for Alcohol Research (A.A.). Funding bodies played no role in data analysis or manuscript preparation.

Contributors

AA, LW, HJE and TF conceived analyses, revised and refined hypotheses and wrote and edited all drafts. AA analyzed data. KKB, JK, SK, MTL and MS provided expertise on phenotypic analyses and clinical significance. HJE, JIN, TF, LW and JT provided expertise on genetic analyses. All authors reviewed the manuscript prior to submission.

Conflicts of interest

LJB is listed as inventor on the patent “Markers for Addiction” (US 20070258898): covering the use of certain SNPs in determining the diagnosis, prognosis, and treatment of addiction and acted as a consultant for Pfizer, Inc. in 2008.

References

- Agrawal, A., Heath, A. C., & Lynskey, M. T. (2011). DSM-IV to DSM-5: The impact of proposed revisions on diagnosis of alcohol use disorders. *Addiction*, 106, 1935–1943.
- Begleiter, H., Reich, T., Hesselbrock, V., Porjesz, B., Li, T. K., Schuckit, M., et al. (1995). The collaborative study on the genetics of alcoholism. *Alcohol Health and Research World*, 19, 228–236.
- Bierut, L. J., Agrawal, A., Bucholz, K. K., Doheny, K. F., Laurie, C., Pugh, E., et al. (2010). A genome-wide association study of alcohol dependence. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 5082–5087.
- Bierut, L. J., Madden, P. A., Breslau, N., Johnson, E. O., Hatsukami, D., Pomerleau, O. F., et al. (2007). Novel genes identified in a high-density genome wide association study for nicotine dependence. *Human Molecular Genetics*, 16, 24–35.

- Bierut, L. J., Strickland, J. R., Thompson, J. R., Afful, S. E., & Cottler, L. B. (2008). Drug use and dependence in cocaine dependent subjects, community-based individuals, and their siblings. *Drug and Alcohol Dependence*, 95, 14–22.
- Blum, K., Liu, Y., Shriner, R., & Gold, M. S. (2011). Reward circuitry dopaminergic activation regulates food and drug craving behavior. *Current Pharmaceutical Design*, 17, 1158–1167.
- Bohn, M. J., Krahn, D. D., & Staehler, B. A. (1995). Development and initial validation of a measure of drinking urges in abstinent alcoholics. *Alcoholism, Clinical and Experimental Research*, 19, 600–606.
- Bonsch, D., Greifenberg, V., Bayerlein, K., Biermann, T., Reulbach, U., Hillemacher, T., et al. (2005). Alpha-synuclein protein levels are increased in alcoholic patients and are linked to craving. *Alcoholism, Clinical and Experimental Research*, 29, 763–765.
- Bonsch, D., Lederer, T., Reulbach, U., Hothorn, T., Kornhuber, J., & Bleich, S. (2005). Joint analysis of the NACP-REP1 marker within the alpha synuclein gene concludes association with alcohol dependence. *Human Molecular Genetics*, 14, 967–971.
- Bonsch, D., Reulbach, U., Bayerlein, K., Hillemacher, T., Kornhuber, J., & Bleich, S. (2004). Elevated alpha synuclein mRNA levels are associated with craving in patients with alcoholism. *Biological Psychiatry*, 56, 984–986.
- Bucholz, K. K., Cadoret, R. J., Cloninger, R. C., Dinwiddie, S. H., Hesselbrock, V., Nurnberger, J. I., et al. (1994). A new, semi-structured psychiatric interview for use in genetic linkage studies. *Journal of Studies on Alcohol*, 55, 149–158.
- Bucholz, K. K., Heath, A. C., Reich, T., Hesselbrock, V. M., Kramer, J. R., Nurnberger, J. I., Jr., et al. (1996). Can we subtype alcoholism? A latent class analysis of data from relatives of alcoholics in a multicenter family study of alcoholism. *Alcoholism, Clinical and Experimental Research*, 20, 1462–1471.
- Buckholtz, J. W., Treadway, M. T., Cowan, R. L., Woodward, N. D., Li, R., Ansari, M. S., et al. (2010). Dopaminergic network differences in human impulsivity. *Science*, 329, 532.
- Carter, B. L., & Tiffany, S. T. (1999). Meta-analysis of cue-reactivity in addiction research. *Addiction*, 94, 327–340.
- Cherpitel, C. J., Borges, G., Ye, Y., Bond, J., Cremonese, M., Moskalewicz, J., et al. (2010). Performance of a craving criterion in DSM alcohol use disorders. *Journal of Studies on Alcohol and Drugs*, 71, 674–684.
- Childress, A. R., Mozley, P. D., McElgin, W., Fitzgerald, J., Reivich, M., & O'Brien, C. P. (1999). Limbic activation during cue-induced cocaine craving. *The American Journal of Psychiatry*, 156, 11–18.
- Diana, M. (2011). The dopamine hypothesis of drug addiction and its potential therapeutic value. *Front Psychiatry*, 2, 64.
- Drummond, D. C. (2001). Theories of drug craving, ancient and modern. *Addiction*, 96, 33–46.
- Drummond, D. C., Litten, R. Z., Lowman, C., & Hunt, W. A. (2000). Craving research: Future directions. *Addiction*, 95(Suppl. 2), S247–S255.
- Ehlers, C. L., & Wilhelmsen, K. C. (2005). Genomic scan for alcohol craving in mission Indians. *Psychiatric Genetics*, 15, 71–75.
- Flannery, B. A., Roberts, A. J., Cooney, N., Swift, R. M., Anton, R. F., & Rohsenow, D. J. (2001). The role of craving in alcohol use, dependence, and treatment. *Alcoholism, Clinical and Experimental Research*, 25, 299–308.
- Foroud, T., Bucholz, K. K., Edenberg, H. J., Goate, A., Neuman, R. J., Porjesz, B., et al. (1998). Linkage of an alcoholism-related severity phenotype to chromosome 16. *Alcoholism, Clinical and Experimental Research*, 22, 2035–2042.
- Foroud, T., Wetherill, L. F., Liang, T., Dick, D. M., Hesselbrock, V., Kramer, J., et al. (2007). Association of alcohol craving with alpha-synuclein (SNCA). *Alcoholism, Clinical and Experimental Research*, 31, 537–545.
- Heinz, A., Siessmeier, T., Wrase, J., Buchholz, H. G., Gruner, G., Kumakura, Y., et al. (2005). Correlation of alcohol craving with striatal dopamine synthesis capacity and D2/3 receptor availability: A combined [¹⁸F]DOPA and [¹⁸F]DMFP PET study in detoxified alcoholic patients. *The American Journal of Psychiatry*, 162, 1515–1520.
- Heinz, A., Siessmeier, T., Wrase, J., Hermann, D., Klein, S., Grusser, S. M., et al. (2004). Correlation between dopamine D(2) receptors in the ventral striatum and central processing of alcohol cues and craving. *The American Journal of Psychiatry*, 161, 1783–1789.
- Hendershot, C. S., Witkiewitz, K., George, W. H., & Marlatt, G. A. (2011). Relapse prevention for addictive behaviors. *Substance Abuse Treatment, Prevention, and Policy*, 6, 17.
- Hesselbrock, M., Easton, C., Bucholz, K. K., Schuckit, M., & Hesselbrock, V. (1999). A validity study of the SSAGA—A comparison with the SCAN. *Addiction*, 94, 1361–1370.
- Irvine, G. B., El-Agnaf, O. M., Shankar, G. M., & Walsh, D. M. (2008). Protein aggregation in the brain: The molecular basis for Alzheimer's and Parkinson's diseases. *Molecular Medicine*, 14, 451–464.
- Kalivas, P. W., & O'Brien, C. (2008). Drug addiction as a pathology of staged neuroplasticity. *Neuropsychopharmacology*, 33, 166–180.
- Kalivas, P. W., & Volkow, N. D. (2005). The neural basis of addiction: A pathology of motivation and choice. *The American Journal of Psychiatry*, 162, 1403–1413.
- Kendler, K. S., Kalsi, G., Holmans, P. A., Sanders, A. R., Aggen, S. H., Dick, D. M., et al. (2011). Genomewide association analysis of symptoms of alcohol dependence in the molecular genetics of schizophrenia (MGS2) control sample. *Alcoholism, Clinical and Experimental Research*, 35, 963–975.
- Keyes, K. M., Krueger, R. F., Grant, B. F., & Hasin, D. S. (2010). Alcohol craving and the dimensionality of alcohol disorders. *Psychological Medicine*, 1–12.
- Keyes, K. M., Krueger, R. F., Grant, B. F., & Hasin, D. S. (2011). Alcohol craving and the dimensionality of alcohol disorders. *Psychological Medicine*, 41, 629–640.
- Koyama, S., Brodie, M. S., & Appel, S. B. (2007). Ethanol inhibition of m-current and ethanol-induced direct excitation of ventral tegmental area dopamine neurons. *Journal of Neurophysiology*, 97, 1777–1785.
- Kozlowski, L. T., Mann, R. E., Wilkinson, D. A., & Poulos, C. X. (1989). “Cravings” are ambiguous: Ask about urges or desires. *Addictive Behaviors*, 14, 443–445.
- Kramer, J. R., Chan, G., Hesselbrock, V. M., Kuperman, S., Bucholz, K. K., Edenberg, H. J., et al. (2010). A principal components analysis of the abbreviated Desires for Alcohol Questionnaire (DAQ). *Journal of Studies on Alcohol and Drugs*, 71, 150–155.

- Larimer, M. E., Palmer, R. S., & Marlatt, G. A. (1999). Relapse prevention. An overview of Marlatt's cognitive-behavioral model. *Alcohol Research & Health*, 23, 151–160.
- Laurie, C. C., Doheny, K. F., Mirel, D. B., Pugh, E. W., Bierut, L., Bhangale, T., et al. (2010). Quality control and quality assurance in genotypic data for genome-wide association studies. *Genetic Epidemiology*, 34(6), 591–602.
- Leggio, L., Ray, L. A., Kenna, G. A., & Swift, R. M. (2009). Blood glucose level, alcohol heavy drinking, and alcohol craving during treatment for alcohol dependence: Results from the Combined Pharmacotherapies and Behavioral Interventions for Alcohol Dependence (COMBINE) study. *Alcoholism, Clinical and Experimental Research*, 33, 1539–1544.
- Lowman, C., Hunt, W. A., Litten, R. Z., & Drummond, D. C. (2000). Research perspectives on alcohol craving: An overview. *Addiction*, 95(Suppl. 2), S45–S54.
- Marlatt, G. A., & Gordon, J. R. (1985). *Relapse prevention*. New York: Guilford Press.
- Mewton, L., Slade, T., McBride, O., Grove, R., & Teesson, M. (2011). An evaluation of the proposed DSM-5 alcohol use disorder criteria using Australian national data. *Addiction*, 106(5), 941–950.
- Monti, P. M., Abrams, D. B., Binkoff, J. A., Zwick, W. R., Liepmann, M. R., Nirenberg, T. D., et al. (1990). Communication skills training, communication skills training with family and cognitive behavioral mood management training for alcoholics. *Journal of Studies on Alcohol*, 51, 263–270.
- Moss, H. B. (2011). Does research support “craving” as a core symptom of substance use disorders in DSM-5? *Psychiatric times* <http://www.psychiatrytimes.com/substance-abuse/content/article/10168/1775007>
- Muthen, L. K., & Muthen, B. O. (2007). *Mplus user's guide* (Fifth ed.). Los Angeles, CA: Muthen & Muthen.
- Narita, M., Soma, M., Tamaki, H., Narita, M., & Suzuki, T. (2002). Intensification of the development of ethanol dependence in mice lacking dopamine D(3) receptor. *Neuroscience Letters*, 324, 129–132.
- O'Brien, C. P. (2005). Anticraving medications for relapse prevention: A possible new class of psychoactive medications. *The American Journal of Psychiatry*, 162, 1423–1431.
- O'Brien, C. (2011). Addiction and dependence in DSM-V. *Addiction*, 106, 866–867.
- Oslin, D. W., Cary, M., Slaymaker, V., Collieran, C., & Blow, F. C. (2009). Daily ratings measures of alcohol craving during an inpatient stay define subtypes of alcohol addiction that predict subsequent risk for resumption of drinking. *Drug and Alcohol Dependence*, 103, 131–136.
- Perez, R. G., Waymire, J. C., Lin, E., Liu, J. J., Guo, F., & Zigmond, M. J. (2002). A role for alpha-synuclein in the regulation of dopamine biosynthesis. *Journal of Neuroscience*, 22, 3090–3099.
- Price, A. L., Patterson, N. J., Plenge, R. M., Weinblatt, M. E., Shadick, N. A., & Reich, D. (2006). Principal components analysis corrects for stratification in genome-wide association studies. *Nature Genetics*, 38, 904–909.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., et al. (2007). PLINK: A tool set for whole-genome association and population-based linkage analyses. *American Journal of Human Genetics*, 81, 559–575.
- SAS Institute (1999). *SAS user guide, version 8.2*. Cary, NC: SAS Institute Inc.
- Sinha, R., Fox, H. C., Hong, K. I., Hansen, J., Tuit, K., & Kreek, M. J. (2011). Effects of adrena sensitivity, stress- and cue-induced craving, and anxiety on subsequent alcohol relapse and treatment outcomes. *Archives of General Psychiatry*, 68, 942–952.
- Sinha, R., Garcia, M., Paliwal, P., Kreek, M. J., & Rounsaville, B. J. (2006). Stress-induced cocaine craving and hypothalamic-pituitary-adrenal responses are predictive of cocaine relapse outcomes. *Archives of General Psychiatry*, 63, 324–331.
- Tupala, E., Hall, H., Bergstrom, K., Sarkioja, T., Rasanen, P., Mantere, T., et al. (2001). Dopamine D(2)/D(3)-receptor and transporter densities in nucleus accumbens and amygdala of type 1 and 2 alcoholics. *Molecular Psychiatry*, 6, 261–267.
- Volkow, N. D., Wang, G. J., Fowler, J. S., Logan, J., Hitzemann, R., Ding, Y. S., et al. (1996). Decreases in dopamine receptors but not in dopamine transporters in alcoholics. *Alcoholism, Clinical and Experimental Research*, 20, 1594–1598.
- Volkow, N. D., Wang, G. J., Telang, F., Fowler, J. S., Logan, J., Childress, A. R., et al. (2006). Cocaine cues and dopamine in dorsal striatum: Mechanism of craving in cocaine addiction. *Journal of Neuroscience*, 26, 6583–6588.
- World Health Organization (2007). *International statistical classification of diseases and related health problems 10th revision* (10th ed.). Geneva, Switzerland.